

Gauge-Symmetric Approach to Effective Lagrangians: The η' meson from QCD

P.H. DAMGAARD
CERN – Geneva

H.B. NIELSEN
The Niels Bohr Institute
Blegdamsvej 17, DK-2100 Copenhagen, Denmark

and

R. SOLLACHER
Gesellschaft für Schwerionenforschung GSI mbH
P.O.Box 110552, D-64220 Darmstadt, Germany

February 1, 2008

Abstract

We present a general scheme for extracting effective degrees of freedom from an underlying fundamental Lagrangian, through a series of well-defined transformations in the functional integral of the cut-off theory. This is done by introducing collective fields in a gauge-symmetric manner. Through appropriate gauge fixings of this symmetry one can remove long-distance degrees of freedom from the underlying theory, replacing them by the collective fields. Applying this technique to QCD, we set out to extract the long-distance dynamics in the pseudoscalar flavour singlet sector through a gauging (and subsequent gauge fixing) of the $U(1)_A$ flavour symmetry which is broken by the anomaly. By this series of exact transformations of a cut-off generating functional for QCD, we arrive at a theory describing the long-distance physics of a pseudoscalar flavour singlet meson coupled to the residual quark-gluon degrees of freedom. As examples of how known low-energy physics can be reproduced in this formulation, we rederive the Witten-Veneziano relation between the η' mass and the topological susceptibility, now for any value of the number of colours N_c . The resulting effective Lagrangian contains an axial vector field, which shares the relevant features with the Veneziano ghost. This field is responsible for removing the η' degree of freedom from the fundamental QCD Lagrangian.

CERN-TH-6959/93
August 1993
hep-th/9308259

1 Introduction

The concept of “effective Lagrangians” has many different meanings, depending on the circumstances under which they have been derived. For example, without knowing the underlying dynamics of certain phenomena in detail, one may try to construct a very general effective field theory consistent with certain global symmetries such as Lorentz invariance, CPT, flavour symmetries, etc. In doing this, it is hoped that only a few of the terms are required to reproduce approximately the observed phenomena, while the lack of knowledge of the underlying theory can be absorbed into a few parameters, – the coupling constants of the effective Lagrangian. Fitting these couplings to experimental observations, one has constructed a well-defined effective theory with predictive power. Chiral Lagrangians in general and linear sigma models are both prime examples of this approach (see, e.g., ref.[1] for an excellent early review). Long before the advent of QCD, these theories accurately described a host of strong-interaction phenomena. Their sole basis was the inferred spontaneously broken global $SU(N_f)_L \times SU(N_f)_R$ chiral symmetry and related basic features of PCAC and current algebra. Clearly, since these models were so successful in describing strong-interaction physics in the low-energy regime, the fine details of the underlying gauged $SU(3)$ theory of quarks and gluons were not very important for these phenomena. A beautiful description of chiral Lagrangians in the light of the more recent developments can be found in a series of papers by Gasser and Leutwyler [2]. These papers also contain a good introduction to the use of chiral perturbation theory.¹

Today, we are of course in a different position with regards to strong-interaction physics. We *know* the underlying theory – QCD – at least down to distance scales that are very far removed from hadronic phenomena. The purpose of an effective field theory for strong interactions is then quite different. Instead of having to solve the underlying theory exactly in the low-momentum regime (something which in principle can be done, at least numerically), we may wish instead to extract directly from the QCD Lagrangian those degrees of freedom that are responsible for the long-distance dynamics. We do not want this in terms of very complicated non-local objects (involving, *e.g.*, the full wave functions of multiplets of bound states), but rather in terms of a preferably local relativistic quantum field theory that to any given level of accuracy will describe the low-energy phenomena. Clearly, this entails extracting from the fundamental Lagrangian those collective degrees of freedom that represent the spectrum of lowest-lying hadrons, and their interactions. In analogy with the usual notion of collective coordinates, we shall view such hadronic excitations as *collective fields* of QCD.

In general, the problem is then the following. Given a local relativistic quantum field theory described in terms of a bare Lagrangian valid up to a cut-off Λ , how do we conveniently extract the relevant physics of an energy scale $M_0 \ll \Lambda$ very far from this ultraviolet cut-off? In other words, which series of manipulations in the path integral will leave us with a Lagrangian that to any degree of required accuracy describes the physics at scale M_0 ? This can be seen as the modern viewpoint on effective Lagrangians of more fundamental theories. It is a viewpoint in which one either explicitly integrates, or at least mentally *imagines* integrating out all degrees of freedom between M_0 and Λ . In this sense, it is a direct application of Wilson’s renormalization group ideas (see, *e.g.*, ref. [4]). Weinberg’s phenomenological Lagrangians [5], and Georgi’s effective Lagrangian scheme [6, 7] are very closely related to this point of view. For two nice expositions of the renormalization group

¹For more recent reviews, and applications to anomalous processes, see *e.g.* ref. [3].

aspects of this approach, see also ref. [8]. The price one pays for integrating out short-distance degrees of freedom is that the full “basis” of field operators appears to a higher and higher extent in the low-energy Lagrangian. This, however, is in any case to be expected of an effective theory. But an obvious drawback of only integrating out short-distance degrees of freedom is that it does not, at least in its most simple formulations, lead to an effective change of variables from, say, quarks and gluons to mesons, baryons and glueballs. The original fermions and vector bosons get more and more “dressed”, but their rôle as constituents of more convenient field variables does not automatically appear.

Indeed, when it comes to the practical question of actually *deriving* directly from QCD an effective long-distance theory, at least one other approach has proven to be more successful. We are referring here to the series of papers in which an effective chiral Lagrangian – describing only those low-energy degrees of freedom that are sensitive to chiral symmetry – has been extracted from the transformation properties of certain generating functionals of QCD [9]. In a more recent paper by Espriu, de Rafael and Taron [10] this idea is beautifully laid out, and carried through to higher orders in both a derivative and $1/N_c$ expansion, N_c being the number of colours. Here, for the first time, an attempt has been made to derive a low-energy chiral Lagrangian directly from QCD.² The agreement with experiment is impressive, in particular when gluonic corrections within the $1/N_c$ expansion are included. At no point has high-momentum modes of the QCD Lagrangian explicitly been integrated out. Instead, such a derivation concerns the “bare” Lagrangians: QCD and the corresponding meson theory with explicit cut-offs.

With these different approaches to the derivation of effective Lagrangians available, we should clearly not introduce yet another method if we cannot significantly shed new light on the problem. We believe that the technique we shall discuss in this paper stands up to this criterion. However, the scheme is not free of difficulties, as we shall describe in detail below. These difficulties are however not inherent in the – as we call it – collective field method itself, but rather stem from the lack of well-defined non-perturbative renormalization prescriptions for continuum formulations of the underlying field theory.

The simplest way to describe our approach is to say that it provides a systematic framework for extracting appropriate collective degrees of freedom (our collective fields), starting directly from the underlying Lagrangian. As we have indicated above, in a renormalization group approach the notion of collective degrees of freedom does not naturally appear³. Yet these degrees of freedom are certainly there (and even becoming increasingly important as one goes to larger distances), and it would be useful to have a method with which to extract them.

A new local gauge symmetry plays a central rôle in our method. This gauge symmetry depends on the change of variables we wish to make, *i.e.* implicitly on the global symmetries we choose our collective fields to probe. It should not be confused with possible local gauge symmetries already inherent in the starting Lagrangian (say, $SU(N_c)$, or $SU(3)$ for QCD), nor should it be given any physical interpretation. It is a technical device introduced only in order to perform certain intermediate steps. In the end, this gauge symmetry is of course completely gauge-fixed. To demystify the meaning of such a new local symmetry, suffice

²For related schemes, see also the contributions in ref. [11].

³Although of course one may choose to introduce such degrees of freedom by hand through the projection operator that links the starting Lagrangian to the “renormalized” Lagrangian which is obtained after integrating out part of the high-momentum modes. With such a choice, we believe one can obtain an effective Lagrangian analogous to the one we shall derive here.

it to say that it can be considered simply as the vehicle with which we can, in a simple and systematic manner, perform complicated changes of variables in the functional integral. Some related considerations can be found in refs.[12, 13]. The method has recently been tested in the solvable realm of two-dimensional field theories where it leads to the standard bosonization rules – plus in fact much more [14, 15]. These exactly solvable two-dimensional examples are very useful for showing the machinery at work in cases where we know the final answer, and we shall return to them below.

In order to extract the collective degrees of freedom describing a flavour singlet pseudoscalar, we are almost unavoidably led to gauging the $U(1)_A$ symmetry. As is well known, this symmetry is anomalous at the quantum level, but this is of no concern here. Our chiral gauge symmetry remains by construction *unbroken*, even if the original global symmetry is broken. The collective field we shall extract from the $U(1)_A$ transformation is related to the η' meson in a manner to be described below.

These general remarks should suffice as an introduction to our approach, and after having outlined the content of the rest of this paper, we shall proceed with the details. In section 2, we present our effective Lagrangian scheme in a situation where the exact answer is known (and an effective Lagrangian approach hence unnecessary), namely the Schwinger model. The obvious analogies to QCD are made explicit by a comparison of chiral Ward identities. Our effective Lagrangian here shows the mass generation of the bound state of a fermion-antifermion pair. In addition, it gives what we interpret as a Lagrangian analogue of the “Veneziano ghost”, which plays such a crucial rôle in the resolution of the 4-dimensional $U(1)$ problem [17]. In section 3, we apply exactly the same procedure to 4-dimensional massless QCD. Despite many formal analogies, the situation here is of course far more complicated, in particular with respect to such aspects as the rôle of the ultraviolet cut-off. However, within the cut-off theory, we can give a reinterpretation of the Witten-Veneziano relation [16, 17] between the η' mass in the limit $N_c \rightarrow \infty$ and the $SU(N_c)$ pure gauge theory topological susceptibility. We discuss the extent to which a similar relation can be derived for any finite value of N_c . At intermediate steps we show how highly non-trivial aspects of the BRST gauge-fixing of the $U(1)_A$ symmetry conspire to leave a very simple final result. As a byproduct, we find again a Lagrangian analogue of the Veneziano ghost, which can be responsible for the saturation of the relevant chiral Ward identities [17]. We briefly comment on the possibility of defining cut-off independent relations of the same kind through an appropriate renormalization procedure. We also discuss some of the general difficulties that are bound to arise in an approach of this kind.

2 A toy model: An effective Lagrangian for the Schwinger model

To illustrate the basic ingredients of the gauge-symmetric approach to effective Lagrangians, we shall start with a simple 2-dimensional toy model: the Schwinger model. This model is solvable in many ways (for a partial list of relevant references, see [18]), most notably in terms of bosonization. We shall here show how the same information can be arrived at through the introduction of a certain collective field $\theta(x)$ without directly bosonizing the theory. The fact that this is possible is important, since we of course have no hope of fully bosonizing 4-dimensional QCD.

Massless QCD and the Schwinger model have many common features. Both theories

show confinement. Usually, they are defined as an ensemble of systems carrying integer instanton number, thus exhibiting a non-trivial vacuum structure. At least for the Schwinger model, this is a necessary ingredient in order to have well-defined asymptotic states [19]; in particular, this guarantees the cluster decomposition property for (vector) gauge-invariant objects. The most interesting aspect with respect to the topics we are discussing in the following is the chiral anomaly in the $U(1)$ sector. We shall list some of the relevant chiral Ward identities for the Schwinger model here, and then compare in the next section with the corresponding 4-dimensional QCD analogues.

We are considering the partition function of the one-flavour Schwinger model

$$\mathcal{Z}_{Schw} = \int \mathcal{D}[\psi, \bar{\psi}, A_\mu] e^{-\int d^2x \mathcal{L}_{Schw}(x)} , \quad (1)$$

where we have put all (vector) gauge-fixing terms into the measure. The Lagrangian in Euclidean space-time reads

$$\mathcal{L}_{Schw} = \frac{1}{4e^2} F_{\mu\nu} F_{\mu\nu} + \bar{\psi}(\not{\partial} - i\not{A})\psi , \quad (2)$$

with the (Euclidean space) conventions

$$\gamma_\mu = \gamma_\mu^\dagger , \quad \{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu} \quad (3)$$

and

$$\gamma_5 = i\gamma_1\gamma_2 = \gamma_5^\dagger , \quad \{\gamma_\mu, \gamma_5\} = 0 . \quad (4)$$

As mentioned at the beginning of this section, we actually have to consider an ensemble of systems with different instanton number. In particular this means that the measure for the vector gauge field is given by

$$\int \mathcal{D}[A_\mu] = \sum_{n=-\infty}^{+\infty} \int \mathcal{D}[A_\mu]_n \quad (5)$$

where, for each n , the vector field A_μ is subject to the constraint

$$\frac{1}{4\pi} \int d^2x \epsilon_{\mu\nu} F_{\mu\nu} = n \quad (6)$$

denoting the instanton number.

Relevant anomalous Ward identities whose derivation will be demonstrated later are

$$0 = \langle \partial_\mu J_\mu^5 + \frac{1}{\pi} \epsilon_{\mu\nu} \partial_\mu A_\nu \rangle , \quad J_\mu^5 = \bar{\psi} \gamma_\mu \gamma_5 \psi , \quad (7)$$

the “anomaly equation”, and

$$\langle \partial_\mu (J_\mu^5(x) + K_\mu(x)) \partial_\mu (J_\mu^5(y) + K_\mu(y)) \rangle = \frac{1}{\pi} \partial^2 \delta(x - y) , \quad (8)$$

the corresponding identity for the correlation function; here K_μ is defined as the topological current of the photon field,

$$K_\mu = \frac{1}{\pi} \epsilon_{\mu\nu} A_\nu . \quad (9)$$

Eqs. (7) and (8) indicate two important properties of the Schwinger model: First of all, there is a conserved axial current $J_\mu^5 + K_\mu$; secondly, there is an associated Goldstone boson. The latter can be seen by introducing an interpolating pseudoscalar field $J_\mu^5 + K_\mu = -if\partial_\mu\varphi$ in eq. (8); this implies a Green's function for $\varphi(x)$ typical of a massless pseudoscalar field and a “decay constant” $f = 1/\sqrt{\pi}$. The latter being independent of the detailed dynamics it may be misleading to call this phenomenon “spontaneous chiral symmetry breaking”. Another common feature of QCD and the Schwinger model is the absence of this Goldstone boson in the physical spectrum. It is widely accepted that this is due to the fact that the conserved axial current, in particular K_μ , is not gauge invariant (see, *e.g.*, [20]).

This model is solved immediately if one applies the usual 2-dimensional bosonization rules, viz.,

$$\bar{\psi}\phi\psi = \frac{1}{2}\partial_\mu\sigma\partial_\mu\sigma \quad , \quad \bar{\psi}\gamma_\mu\psi = \frac{1}{\sqrt{\pi}}\epsilon_{\mu\nu}\partial_\nu\sigma \quad , \quad (10)$$

which allows us to rewrite the Lagrangian in the form

$$\mathcal{L}_{Schw} = \frac{1}{2}\partial_\mu\sigma\partial_\mu\sigma + \frac{i}{\sqrt{\pi}}\partial_\mu\sigma \epsilon_{\mu\nu}A_\nu + \frac{1}{4e^2}F_{\mu\nu}F_{\mu\nu} \quad . \quad (11)$$

This Lagrangian can be diagonalized in a shortcut manner if one uses the electric field strength F_{12} instead of the vector field A_μ :

$$\mathcal{L}_{Schw} = \frac{1}{2}\partial_\mu\sigma\partial_\mu\sigma - \frac{i}{\sqrt{\pi}}\sigma F_{12} + \frac{1}{2e^2}F_{12}^2 \quad . \quad (12)$$

A simple shift

$$F_{12} = F'_{12} - \frac{ie^2}{\sqrt{\pi}}\sigma \quad (13)$$

yields

$$\mathcal{L}_{Schw} = \frac{1}{2}\partial_\mu\sigma\partial_\mu\sigma + \frac{e^2}{2\pi}\sigma^2 + \frac{1}{2e^2}F_{12}^{\prime 2} \quad . \quad (14)$$

From this we read off that σ describes a free massive pseudoscalar of mass $m = e/\sqrt{\pi}$. From the bosonization rules, we note that it can be regarded as a $\bar{\psi}\psi$ bound state.

Let us now consider the electric field strength F_{12} in more detail. For its Green's function we find

$$\begin{aligned} \langle F_{12}(x)F_{12}(y) \rangle &= \left\langle \left(F'_{12}(x) - \frac{ie^2}{\sqrt{\pi}}\sigma(x) \right) \left(F'_{12}(y) - \frac{ie^2}{\sqrt{\pi}}\sigma(y) \right) \right\rangle \\ &= \langle F'_{12}(x)F'_{12}(y) \rangle - \frac{e^4}{\pi}\langle \sigma(x)\sigma(y) \rangle \end{aligned} \quad (15)$$

In (Euclidean) momentum space this reads

$$\begin{aligned} G_F(p) &= G_{F'}(p) - \frac{e^4}{\pi}G_\sigma(p) \\ &= e^2 - \frac{e^4}{\pi} \frac{1}{p^2 + \frac{e^2}{\pi}} \end{aligned} \quad (16)$$

This Green's function is closely related to the topological susceptibility,

$$G_{top} = \int d^2x \left\langle \frac{1}{2\pi}\epsilon_{\mu\nu}F_{\mu\nu}(x) \frac{1}{2\pi}\epsilon_{\mu\nu}F_{\mu\nu}(0) \right\rangle$$

$$\begin{aligned}
&= \frac{1}{\pi^2} G_F(p=0) \\
&= \lim_{p \rightarrow 0} \left(f^2 m^2 - \frac{f^2 m^4}{p^2 + m^2} \right) = 0 \quad ,
\end{aligned} \tag{17}$$

with f and m defined above. This result is the 2-dimensional counterpart of what Witten [16] suggested to be the case for QCD, namely a cancellation between a pure (vector) gauge field and a massive pseudoscalar contribution to the topological susceptibility.

Let us now show how the same result can be obtained from our effective Lagrangian point of view. The first step of the construction consists in performing a suitable field-enlarging transformation in the functional integral (1). Wishing in the process to introduce a flavour-singlet pseudoscalar field, call it $\theta(x)$, an obvious choice is to perform a local chiral rotation of the quark fields with $\theta(x)$ acting as the chiral angle:

$$\psi(x) = e^{i\theta(x)\gamma_5} \chi(x) \quad , \quad \bar{\psi}(x) = \bar{\chi}(x) e^{i\theta(x)\gamma_5}. \tag{18}$$

Since this defines the original quark fields in terms of a local product of new fields, it is essential that the field transformation be made only in a regularized version of the generating functional. In choosing the regulator we also have to be careful: It turns out that for a chiral transformation of the kind (18), an integrability condition can only be satisfied within a restricted class of ultraviolet regulators. This is known as the Wess-Zumino consistency condition (see, *e.g.*, ref. [21]), and a regularization scheme satisfying this requirement is similarly called consistent. A convenient consistent scheme in the fermion sector is provided by a set of Pauli-Villars regulator fields, and this is what we shall be using in this paper.

Having thus defined the corresponding Pauli-Villars regularized generating functional, we can now see the consequences of the field transformation (18) on this generating functional. We will get two pieces, one “classical” stemming from the variation of the action in eq.(1) under the local chiral rotation, and one “quantum mechanical” from the change of the fermionic measure. The former part is obtained trivially. The latter, which if we reinstate factors of Planck’s constant is proportional to $\hbar$, can be computed by means of the method described in [21]. One central property of this quantum mechanical part coming from the fermionic Jacobian is that it can be rearranged into an expansion in decreasing powers of Λ , the ultraviolet cut-off. In this 2-dimensional example, the cut-off can eventually be taken to infinity in a straightforward manner, and we shall therefore only be concerned with the leading terms in this $1/\Lambda$ expansion. (This is one of the crucial differences between the 2-dimensional example and QCD, see section 3 below.)

With these remarks in mind, we can write the chirally rotated Schwinger model in the form

$$\mathcal{L}_{ext} = \frac{1}{4e^2} F_{\mu\nu} F_{\mu\nu} + \bar{\chi}(\not{\partial} - i\not{A} + i\not{\partial}\theta\gamma_5)\chi + \frac{1}{2\pi} \partial_\mu \theta \partial_\mu \theta - \frac{i}{2\pi} \theta \epsilon_{\mu\nu} F_{\mu\nu} . \tag{19}$$

At this stage we are able to derive the Ward identities (7) and (8) from the invariance of the partition function (1) under variable transformations.

We have now completed the part of the field-enlarging transformation. Next, we promote the collective field $\theta(x)$ to a true dynamical field of the theory. In the path integral formulation it means that we integrate over $\theta(x)$ as well. This can be done without changing the generating functional if we note that it corresponds to the introduction of gauge degrees of freedom, and that these degrees of freedom need to be gauge fixed [23]. The corresponding gauge symmetry reads in this case:

$$\chi(x) \rightarrow e^{i\alpha(x)\gamma_5} \chi(x)$$

$$\begin{aligned}\bar{\chi}(x) &\rightarrow \bar{\chi}(x)e^{i\alpha(x)\gamma_5} \\ \theta(x) &\rightarrow \theta(x) - \alpha(x).\end{aligned}\tag{20}$$

We have already pointed out that introducing a functional integration over θ renders the extended partition function with the Lagrangian (19) symmetric under the gauge transformation (20). At first sight, this may appear surprising: After all, the fermionic functional measure is precisely *not* invariant under chiral rotations such as the one of eq. (20). Does this not spoil the gauge symmetry at the quantum level? The answer is no: A gauge symmetry introduced in the way described here *automatically* takes into account any non-trivial transformation of the measure as well. Indeed, the *action* of eq.(19) is not at all invariant under the chiral gauge symmetry of eq.(20). Only if we correctly include the additional terms from the measure is the whole expression invariant. At a more formal level, by switching to an overcomplete basis of fields as indicated, and integrating over all of these, one introduces a set of Hamiltonian constraints. These constraints are the generators of the gauge symmetry [22, 23]. As we have seen, this holds in the case of anomalous symmetries as well! (Note, incidentally, that we are not gauging the chiral symmetry in the usual way by means of spin-1 axial vector fields; the extra gauge degrees of freedom are here carried entirely by the spin-0 pseudoscalar field $\theta(x)$, the derivative of which corresponds to a “pure gauge” in axial-vector gauge theories.) One amusing consequence of this is that if one gauges an anomalous global symmetry in this manner, the resulting *unbroken* Ward identities derived by, *e.g.*, the residual BRST symmetry left after gauge fixing are identical to the *anomalous* Ward identities of the original global symmetry. This holds here as well.

In order to recover the original generating functional, we have to gauge-fix this symmetry. We are obviously free to do this in any manner we want. If, for example, we simply gauge-fix on the trivial surface $\theta(x) = 0$, we just recover the starting Lagrangian. But we are not required to make such a trivial gauge choice. Our only constraint is that we must remove one and only one degree of freedom – the one we artificially added by going to a field basis enlarged by the θ sector. We shall choose to do so by gauge fixing a function involving fermionic bilinears. For that purpose, it is useful to note that the axial current shifts under (18) as (for details see [14])

$$i\bar{\psi}\gamma_\mu\gamma_5\psi = i\bar{\chi}\gamma_\mu\gamma_5\chi - \frac{1}{\pi}\partial_\mu\theta \quad .\tag{21}$$

The whole expression is, of course, gauge invariant; however, the individual components on the r.h.s. are gauge dependent. The last term in eq. (21) is induced by the axial anomaly.

We use this manifestation of the chiral anomaly as a guide to choosing a convenient non-trivial gauge fixing. Consider

$$\Phi = \Delta i\pi \frac{\partial_\mu}{\partial^2} \bar{\chi}\gamma_\mu\gamma_5\chi - (1 - \Delta)\theta \quad ,\tag{22}$$

where Δ is a free parameter. This choice of gauge implies that the θ -dependent part of the divergence of the axial current should describe a fraction Δ of the divergence of the *physical* axial current expressed through the *original* fermionic degrees of freedom $\bar{\psi}$ and ψ .

The above gauge-fixing function is, however, defined in a very formal manner on account of the inverse Laplacian. Furthermore, if we wish to implement Φ as a δ -function constraint in the path integral, we should be careful that it satisfies correct transformation properties. Under a global chiral rotation,

$$\bar{\chi}(x) \rightarrow \bar{\chi}(x)e^{i\alpha\gamma_5} \quad , \quad \chi(x) \rightarrow e^{i\alpha\gamma_5}\chi(x) \quad ,\tag{23}$$

the object involving $\bar{\chi}\gamma_\mu\gamma_5\chi$ remains *classically* invariant but quantum mechanically it shifts owing to the chiral anomaly. We find the shift similar to that for (21):

$$i\pi\frac{\partial_\mu}{\partial^2}\bar{\chi}\gamma_\mu\gamma_5\chi \rightarrow i\pi\frac{\partial_\mu}{\partial^2}\bar{\chi}\gamma_\mu\gamma_5\chi + \alpha . \quad (24)$$

The formal operator ∂^{-2} is included precisely in order to cancel the additional operator multiplying α .

The action of the original Lagrangian $\mathcal{L}_{Schw}$ does not remain invariant under constant chiral rotations, but shifts because of the axial anomaly:

$$S = \int d^2x \mathcal{L}_{Schw} \rightarrow S - \frac{i}{2\pi}\alpha \int d^2x \epsilon_{\mu\nu} F_{\mu\nu} = S - 2in\alpha . \quad (25)$$

We assume again that we sum only over integer winding numbers. The action does, however, remain invariant under constant chiral rotations of the form $\alpha = n'\pi$ with n' another integer. On the other hand, this means that $\theta(x)$ is only globally defined modulo π .

Gauge-fixing the constant $\theta(x)$ modes is then non-trivial. The gauge-fixing constraint must respect the above periodicity property; there must, even in the gauge-fixed functional integral, be no distinction between $\theta(x)$ and $\theta(x) + n'\pi$. If we choose a δ -function constraint to implement the gauge choice, this δ -function must then necessarily be *globally periodic*. When fixing a definite value for $\theta(x)$, as for a gauge-fixing function $\Phi = F(x) - \theta(x)$, this implies

$$\delta(F(x) - \theta(x)) = \delta(F(x) - \theta(x) - n'\pi) . \quad (26)$$

This is exactly what is achieved by Φ above. Under a constant chiral gauge transformation of magnitude $n'\pi$,

$$\Phi \rightarrow \Phi + \Delta n'\pi + (1 - \Delta)n'\pi = \Phi + n'\pi . \quad (27)$$

Thus, if we wish to enforce the gauge choice Φ through a δ -function constraint inside the path integral, this δ -function must be globally periodic with period $n\pi$. We can represent such a globally periodic δ -function by means of a Nakanishi-Lautrup auxiliary field $b(x)$, so that this gauge-fixing function just provides a few new terms in the action.

In practice, this is done by implementing the globally periodic δ -function using a functional Fourier representation:

$$\begin{aligned} \delta(\Phi) &= \int \mathcal{D}[b] \exp \left[+ \int d^2x d^2z b(x) \left(i\pi\Delta\partial_{(x-z)}^{-2}\partial_\mu^{(z)}j_\mu^5(z) - (1 - \Delta)\delta(x - z)\theta(z) \right) \right] \\ &= \int \mathcal{D}[b] \exp \left[+ \int d^2x \left(i\pi\Delta B_\mu(x)j_\mu^5(x) + (1 - \Delta)\theta(x)\partial_\mu B_\mu(x) \right) \right] \end{aligned} \quad (28)$$

Here, $j_\mu^5 = \bar{\chi}\gamma_\mu\gamma_5\chi$ denotes the axial vector current of the transformed fermions; the axial vector field $B_\mu(x)$ is defined by

$$B_\mu(x) \equiv \int d^2y d^2z b(y)\partial_{(y-z)}^{-2}\partial_\mu^{(z)}\delta(z - x) . \quad (29)$$

Note that this implies $b(x) = -\partial_\mu B_\mu(x)$. A globally periodic δ -function can be represented by

$$\delta(\Phi(x)) = \sum_{k=-\infty}^{\infty} \int \mathcal{D}[b]_k \exp \left[\int d^2x b(x)\Phi(x) \right] , \quad (30)$$

where b is constrained:

$$\int d^2x \, b(x) = - \int d^2x \, \partial_\mu B_\mu = 2ik . \quad (31)$$

This global constraint means that $b(x)$ (or $\partial_\mu B_\mu$) share certain properties with topologically non-trivial fields [14].

The gauge-fixing is complete once the appropriate ghost term is found. This can be achieved by requiring BRST symmetry in the gauge-fixed Lagrangian. In contrast with the usual situation, the functional measure is not invariant under the BRST transformations

$$\begin{aligned} \delta \bar{\chi}(x) &= i \bar{\chi}(x) \gamma_5 c(x) \\ \delta \chi(x) &= -i c(x) \gamma_5 \chi(x) \\ \delta \theta(x) &= -c(x) \\ \delta c(x) &= 0 \\ \delta \bar{c}(x) &= b(x) \\ \delta b(x) &= 0 . \end{aligned} \quad (32)$$

This property, however, is responsible for the gauge invariance (and therefore BRST invariance) of the partition function of the modified Lagrangian $\mathcal{L}_{ext}$. The only BRST-variant term is therefore the one that has been added to the modified $\mathcal{L}_{ext}$ by imposing the gauge constraint. Its BRST variation is

$$\delta [b(x)\Phi(x)] = b(x)c(x) . \quad (33)$$

From that we immediately see that we need a trivial ghost term, namely

$$\mathcal{L}_{ghost} = \bar{c}c , \quad (34)$$

whose BRST variation precisely cancels the one above. This is just what one obtains when BRST gauge-fixing in the standard manner by adding a term $\delta[\bar{c}\Phi]$ to the action. Obviously, the ghosts decouple and will be neglected in the following.

The complete gauge fixed Lagrangian now reads

$$\begin{aligned} \mathcal{L}'_{Schw} &= \bar{\chi} \left(\not{\partial} - i\not{A} - i(\pi \Delta \not{B} - \not{\partial} \theta) \gamma_5 \right) \chi + \bar{c}c + \frac{1}{4e^2} F_{\mu\nu} F_{\mu\nu} \\ &\quad + \frac{1}{2\pi} \partial_\mu \theta \partial_\mu \theta - \frac{i}{2\pi} \theta \epsilon_{\mu\nu} F_{\mu\nu} - (1 - \Delta) \theta \partial_\mu B_\mu . \end{aligned} \quad (35)$$

From now on, let us concentrate on the gauge $\Delta = 1$, which in the language of ref. [14] is called the “bosonization gauge”.

The bosonization of the Schwinger model is now a one-step procedure. Shifting B_μ as

$$B_\mu \rightarrow B_\mu + \frac{1}{\pi} \partial_\mu \theta + \frac{i}{\pi} \epsilon_{\mu\nu} A_\nu , \quad (36)$$

thereby taking advantage of the relation $\gamma_\mu \gamma_5 = i \epsilon_{\mu\nu} \gamma_\nu$ for 2-dimensional γ -matrices, the fermions decouple completely from A_μ and θ . The Lagrangian now reads

$$\begin{aligned} \mathcal{L}_{bos} &= \bar{\chi} (\not{\partial} - i\pi \not{B} \gamma_5) \chi + \bar{c}c \\ &\quad + \frac{1}{4e^2} F_{\mu\nu} F_{\mu\nu} + \frac{1}{2\pi} \partial_\mu \theta \partial_\mu \theta - \frac{i}{2\pi} \theta \epsilon_{\mu\nu} F_{\mu\nu} \end{aligned} \quad (37)$$

The fields $\bar{\chi}, \chi, B_\mu, \bar{c}$ and c can now be integrated out; this provides a normalization factor necessary to yield the correct normalization with respect to the original fermionic representation of the partition function (1).

The careful reader may wonder whether we are allowed to perform the shift of B_μ . We recall that in general A_μ can be written as [24]

$$A_\mu(x) = C_\mu(x) + \epsilon_{\mu\nu} \partial_\nu \phi(x) + \partial_\mu \varphi(x) \quad (38)$$

where C_μ carries the topological information, *i.e.*

$$\frac{1}{2\pi} \int d^2x \epsilon_{\mu\nu} \partial_\mu C_\nu = n \quad (39)$$

The term $\partial_\mu \varphi(x)$ is a pure (vector) gauge component; it can be removed either by a corresponding (vector) gauge transformation of the fermion fields or by a suitable (vector) gauge condition. The shift of B_μ then means

$$B_\mu \rightarrow B_\mu + \frac{1}{\pi} \partial_\mu \theta - \frac{i}{\pi} \partial_\mu \phi + \frac{i}{\pi} \epsilon_{\mu\nu} C_\nu \quad (40)$$

This implies

$$- \partial_\mu B_\mu = b \rightarrow b - \frac{1}{\pi} \partial^2 \theta + \frac{i}{\pi} \partial^2 \phi - \frac{i}{\pi} \epsilon_{\mu\nu} \partial_\mu C_\nu \quad (41)$$

and therefore, after integration over euclidean space-time,

$$2ik \rightarrow 2ik - 2in \quad (42)$$

The latter transformation is just a shift of the summation index k by the instanton number n defining a new summation index k' . This is a very important point: only with the gauge fixing with a globally periodic δ -function do we arrive at the bosonized version of the Schwinger model. Any other choice of gauge would presumably *not* lead to a *complete* decoupling of the fermions from the vector field A_μ . Another aspect of this gauge will be discussed in a moment.

We are now presenting a more detailed investigation of the mass generation mechanism starting again from the Lagrangian (35). The idea is to construct an effective Lagrangian for the field θ using as few properties special to two dimensions as possible. For this purpose we take again (35) with $\Delta = 1$ and integrate out all fields except θ . The part of the action containing the coupling of θ to the vector gauge field and to the fermionic axial current, denoted by X in the next equation, is treated in a cumulant expansion,

$$\langle e^X \rangle = e^{\langle X \rangle + \frac{1}{2} \langle X^2 \rangle_c + \dots} \quad (43)$$

where $\langle X^2 \rangle_c = \langle X^2 \rangle - \langle X \rangle^2$ denotes the “connected” part of the corresponding moment. The expectation values $\langle \dots \rangle$ have to be taken with respect to a *truncated* theory where θ is set to zero. In the following, we will indicate this by an index $_{trunc}$. Furthermore, due to the gauge constraint represented by a coupling between B_μ and $\bar{\chi} \gamma_\mu \gamma_5 \chi$ at least the divergence of the axial fermion current is identically zero and therefore we can neglect the coupling of θ to the axial fermion current; this effect of the gauge constraint could also be seen by shifting B_μ by $\partial_\mu \theta / \pi$.

This leads to an effective action

$$S_{eff} = \int d^2x \left(\frac{1}{2\pi} \partial_\mu \theta(x) \partial_\mu \theta(x) - \frac{i}{2\pi} \theta(x) \langle \epsilon_{\mu\nu} F_{\mu\nu}(x) \rangle_{trunc} \right)$$

$$\begin{aligned}
& + \int d^2x \int d^2y \frac{1}{2} \theta(x) \left\langle \frac{1}{2\pi} \epsilon_{\mu\nu} F_{\mu\nu}(x) \frac{1}{2\pi} \epsilon_{\mu\nu} F_{\mu\nu}(y) \right\rangle_{trunc,c} \\
& + \dots
\end{aligned} \tag{44}$$

As mentioned, the expectation values $\langle \dots \rangle_{trunc}$ are to be taken with respect to the truncated system described by the Lagrangian

$$\mathcal{L}_{trunc} = \frac{1}{4e^2} F_{\mu\nu} F_{\mu\nu} + \bar{\chi} (\not{\partial} - i\not{A} - i\pi \not{B} \gamma_5) \chi + \bar{c} c \quad . \tag{45}$$

The manipulations leading to (44) and (45) did not involve any special properties of 2 dimensional field theory and should be extendable to higher dimensions. This will be demonstrated in the next section.

In order to investigate the effective Lagrangian further, we now have to make use of known results in two dimensions. The easiest way to solve the truncated theory (45) is to shift B_μ by $i\epsilon_{\mu\nu} A_\nu / \pi$ as before. This decouples the fermions and the (vector) gauge field A_μ . Another point of view of this decoupling was presented in connection with bosonization of massive theories in two dimensions [14]; the field $\epsilon_{\mu\nu} B_\nu$ is essentially the same as a (vector) gauge field like A_μ , up to pure (vector) gauge degrees of freedom. The difference to the field A_μ is the lack of a kinetic term. This can be interpreted as an infinitely strong coupling. In two dimensions this means total confinement; all (physical) correlation functions are zero. The only non-vanishing expectation value is the scalar condensate $\langle \bar{\chi} \chi \rangle_{trunc}$. This implies the vanishing of the coupling between A_μ and the fermions.

What remains for the (vector) gauge field A_μ is the kinetic term alone. This implies a correlation function

$$\left\langle \frac{1}{2\pi} \epsilon_{\mu\nu} F_{\mu\nu}(x) \frac{1}{2\pi} \epsilon_{\mu\nu} F_{\mu\nu}(y) \right\rangle_{trunc,c} = \frac{e^2}{\pi^2} \delta(x-y) \quad ; \tag{46}$$

all higher correlation functions are vanishing. As a consequence, the expansion in (44) terminates at second order and we are thus back at our diagonalized version (14). The cumulant expansion leads in this case to the exact result!

We now concentrate on the field B_μ . By simple shifts we get the following Ward identity for the truncated theory:

$$\langle \partial_\mu (j_\mu^5(x) + K_\mu(x) + iB_\mu(x)) \rangle = 0 \quad . \tag{47}$$

Recall that (45) looks almost like the Lagrangian of the original Schwinger model, except for the coupling to B_μ . In the truncated theory the current $j_\mu^5 + K_\mu$ is no longer conserved due to the presence of precisely this topologically non-trivial field B_μ .

The consequences of this broken chiral symmetry are quite drastic. Consider another Ward identity

$$\langle \partial_\mu B_\mu(x) \partial_\nu B_\nu(y) \rangle_{trunc,c} = \frac{1}{\pi} \partial^2 \delta(x-y) - \left\langle \frac{1}{2\pi} \epsilon_{\mu\nu} F_{\mu\nu}(x) \frac{1}{2\pi} \epsilon_{\mu\nu} F_{\mu\nu}(y) \right\rangle_{trunc,c} \quad . \tag{48}$$

Using (46) and recalling the purely longitudinal character of B_μ , its 2-point function in momentum space reads

$$\int d^2x e^{-ipx} \langle B_\mu(x) B_\nu(0) \rangle_{trunc,c} = -f^2 \frac{p_\mu p_\nu}{p^4} (p^2 + m^2) \quad , \tag{49}$$

implying a behaviour like $p_\mu p_\nu / p^4$ in the limit $p^2 \rightarrow 0$. This behaviour can also be deduced from (48) after integrating over x :

$$\int d^2x \langle \partial_\mu B_\mu(x) \partial_\nu B_\nu(y) \rangle_{trunc} = - \int d^2x \left\langle \frac{1}{2\pi} \epsilon_{\mu\nu} F_{\mu\nu}(x) \frac{1}{2\pi} \epsilon_{\mu\nu} F_{\mu\nu}(y) \right\rangle_{trunc} \quad (50)$$

This equation relates the low-momentum behaviour of the 2-point function of B_μ to the topological susceptibility of the truncated system. We already know that the latter is different from zero; this implies a $p_\mu p_\nu / p^4$ -behaviour in the limit $p^2 \rightarrow 0$. Thus, from very general arguments we find that B_μ has all the characteristic features of the “axial four-vector ghost” (here axial two-vector ghost) suggested by Veneziano [17]. We believe that one may regard B_μ as one explicit realization of this ghost.

We would like to emphasize that the discussion of the Veneziano ghost was based mainly on the Ward identity (50), which can be derived from chiral transformations and shifts of B_μ . This suggests a possible generalization to four dimensions. Indeed, we shall see that the discussion concerning mass generation and the Veneziano ghost can be formally repeated in four dimensions with the same arguments as in the 2-dimensional case. However, it will turn out that there are also major differences due to the much more complicated dynamics of QCD.

3 A flavour-singlet pseudoscalar meson from QCD

The simplest setting with which to illustrate how our method works in the case of QCD, is that of describing the long-distance dynamics of the η' meson. Because it is a flavour singlet, the more complicated machinery of non-Abelian chiral transformations is not needed in this case. We can then describe our method in simple terms, while still having a physical situation in mind. The only complication in this case arises from the chiral anomaly, but also this can be taken fully into account, just as in the previous 2-dimensional example. The flavour-singlet states are by no means the lightest long-distance excitations of QCD, but as shall become clear in the following, this poses no problems either. All dynamics of the lighter bound states will in this case still be contained in the (rotated) quark-gluon sector.

Our starting point is in this case a generating functional for QCD (in Euclidean space) of the form

$$\begin{aligned} \mathcal{Z}_{QCD}[V, A] &= \int \mathcal{D}[\bar{\psi}, \psi] \mathcal{D}\mu[G] e^{-\int d^4x \mathcal{L}_{QCD}} \\ \mathcal{L}_{QCD} &= \bar{\psi}(x) (\not{\partial} - i\not{G}(x) - i\not{M}(x) - i\not{A}(x)\gamma_5) \psi(x) + \frac{1}{4g^2} \text{tr} G_{\mu\nu}(x) G_{\mu\nu}(x). \end{aligned} \quad (51)$$

Here $V_\mu(x)$ is an external vector source and $A_\mu(x)$ an external axial vector source, both Abelian (diagonal in the $SU(N_f)$ flavour indices). The vector potential $G_\mu(x)$ is the usual gluon field, here for convenience generalized to $SU(N_c)$, and $G_{\mu\nu}(x)$ is the corresponding field strength tensor:

$$G_{\mu\nu} = \partial_\mu G_\nu - \partial_\nu G_\mu - i[G_\mu, G_\nu]. \quad (52)$$

The usual $SU(N_c)$ colour gauge symmetry of course has to be gauge-fixed in the standard manner, including also Yang-Mills ghosts. For the moment, we simply include these Yang-Mills gauge-fixing terms implicitly in the gluon measure $\mathcal{D}\mu[G]$.

There is nothing unphysical implied by the coupling to external vector and axial vector sources; these sources only serve to define appropriate Green functions through functional

differentiation. They are clearly not intrinsically part of QCD, and will eventually be set equal to zero. Nevertheless, they turn out to play a rather profound rôle in the derivation of the effective Lagrangian. They would also, of course, acquire a physical meaning if they were to include the couplings of the electroweak interactions.

The γ -matrices are Hermitean and obey the algebra

$$\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu} \quad , \quad \{\gamma_5, \gamma_\mu\} = 0 \quad , \quad \gamma_5 = -\gamma_1\gamma_2\gamma_3\gamma_4 \quad . \quad (53)$$

As in the 2-dimensional example, it is essential that the field transformation is made only in a regularized version of the QCD generating functional. A convenient consistent scheme in the fermion sector is again provided by a set of Pauli-Villars regulator fields. It should be kept in mind that these regulators *only* regularize the fermionic sector of QCD. Although we shall not consider field transformations involving the colour gauge potentials, we of course still need to regularize also the gluon sector of QCD. To interpret the resulting field-transformed Lagrangian as an effective Lagrangian with an ultraviolet cut-off Λ , a regularization procedure with a similar cut-off ought to be introduced in the gluonic sector. This is a non-trivial issue, but at least one correct cut-off procedure is known to exist [28].

Upon the field transformation we will get two pieces, one classical from the variation of the action in eq.(51) under the local chiral rotation, and one quantum mechanical from the change of the fermionic measure. The latter part can again essentially be read off from ref. [21]. We rearrange it into an expansion in decreasing powers of Λ , the ultraviolet cut-off. Explicitly, we can then write the regularized generating functional as

$$\begin{aligned} \mathcal{Z}_\Lambda[V, A] &= \int \mathcal{D}_\Lambda[\bar{\chi}, \chi] \mathcal{D}\mu[G] e^{-\int d^4x \mathcal{L}'} \\ \mathcal{L}' &= \frac{1}{4g^2} \text{tr} G_{\mu\nu} G_{\mu\nu} + \bar{\chi}(\not{\partial} - i\not{A} - i\not{A}\gamma_5 + i\not{\partial}\theta\gamma_5)\chi \\ &\quad + \mathcal{L}_{WZ} + \mathcal{L}_J \end{aligned} \quad (54)$$

where the last two terms arise from the Jacobian of the transformation. As advertized, these last two terms can be computed to all orders in an expansion in decreasing powers of the ultraviolet cut-off Λ . The first part, the Wess-Zumino term, starts at $\mathcal{O}(\Lambda^0)$:

$$\mathcal{L}_{WZ} = -\frac{iN_f}{16\pi^2} \theta \epsilon_{\mu\nu\rho\sigma} \left(\text{tr} G_{\mu\nu} G_{\rho\sigma} - 4N_c \partial_\mu V_\nu \partial_\rho V_\sigma - \frac{4N_c}{3} \partial_\mu A_\nu \partial_\rho A_\sigma \right) + \mathcal{O}(\Lambda^{-2}), \quad (55)$$

while the second part – the positive-parity part, in contrast with the negative-parity Wess-Zumino term – starts at $\mathcal{O}(\Lambda^2)$:

$$\mathcal{L}_J = \Lambda^2 \mathcal{L}_2 + \mathcal{L}_0 + \frac{1}{\Lambda^2} \mathcal{L}_{-2} + \frac{1}{\Lambda^4} \mathcal{L}_{-4} + \dots \quad (56)$$

where

$$\begin{aligned} \mathcal{L}_2 &= \frac{N_f N_c \kappa_2}{4\pi^2} \left(A_\mu A_\mu - (A_\mu - \partial_\mu \theta)(A_\mu - \partial_\mu \theta) \right) \\ \mathcal{L}_0 &= \frac{N_f N_c}{24\pi^2} \left(\partial_\mu A_\nu \partial_\mu A_\nu - \partial_\mu (A_\nu - \partial_\nu \theta) \partial_\mu (A_\nu - \partial_\nu \theta) \right. \\ &\quad \left. + 2(A_\mu A_\mu)^2 - 2((A_\mu - \partial_\mu \theta)(A_\mu - \partial_\mu \theta))^2 \right) \\ \mathcal{L}_{-2} &= \frac{N_f \kappa_{-2}}{48\pi^2} \left(N_c \partial^2 A_\mu \partial^2 A_\mu - N_c \partial^2 (A_\mu - \partial_\mu \theta) \partial^2 (A_\mu - \partial_\mu \theta) \right) \end{aligned}$$

$$+\left(A_\mu A_\mu - (A_\mu - \partial_\mu \theta)(A_\mu - \partial_\mu \theta)\right) \text{tr}_c G_{\nu\rho} G_{\nu\rho} + \mathcal{O}(A_\mu^4) \Big) \quad (57)$$

Here, tr_c denotes a trace over colour indices. In (57) we list only the first three terms; the whole expansion in increasing powers of inverse cut-off can be computed following the technique described in ref. [21]. Similarly, we show explicitly in (55) only the leading contribution.

The coefficients κ_2 and κ_{-2} in eq. (57) are regularization-scheme dependent constants, given here by

$$\kappa_n = \sum_i c_i k_i^n \ln k_i^2 \quad , \quad \sum_i c_i = 1 \quad , \quad \sum_i c_i k_i^m = 0 \quad \text{for} \quad m = 1, \dots, 4 \quad (58)$$

where the k_i 's are the Pauli-Villars regulator masses in units of the cut-off Λ , *i.e.* $M_i = k_i \Lambda$.

This completes the part of the field-enlarging transformation in QCD. When we next integrate over the collective fields in the path integral, a chiral gauge symmetry again appears:

$$\begin{aligned} \chi(x) &\rightarrow e^{i\alpha(x)\gamma_5} \chi(x) \\ \bar{\chi}(x) &\rightarrow \bar{\chi}(x) e^{i\alpha(x)\gamma_5} \\ \theta(x) &\rightarrow \theta(x) - \alpha(x). \end{aligned} \quad (59)$$

We shall return to the question of gauge fixing in section 3.2.

One important property of the transformed action in (54) is that it already contains a kinetic energy piece for the collective field $\theta(x)$. We can write the full Lagrangian in (54) in a more conventional form of a pseudoscalar field coupled to the rotated fermion fields, the gluons and the external sources:

$$\begin{aligned} \mathcal{L}' = & \frac{1}{4g^2} \text{tr} G_{\mu\nu} G_{\mu\nu} + \bar{\chi} (\not{\partial} - i\not{G} - i\not{M} - i\not{A}\gamma_5 + i\not{\partial}\theta\gamma_5) \chi \\ & + \frac{N_f}{2} \partial_\mu \theta f^2 \partial_\mu \theta - N_f A_\mu f^2 \partial_\mu \theta \\ & - \frac{N_f N_c}{12\pi^2} \left(\left((A_\mu - \partial_\mu \theta)(A_\mu - \partial_\mu \theta) \right)^2 - (A_\mu A_\mu)^2 \right) \\ & - \theta \frac{iN_f}{16\pi^2} \epsilon_{\mu\nu\rho\sigma} \left(\text{tr}_c G_{\mu\nu} G_{\rho\sigma} - 4N_c \partial_\mu V_\nu \partial_\rho V_\sigma - \frac{4N_c}{3} \partial_\mu A_\nu \partial_\rho A_\sigma \right) \\ & + \mathcal{O}(\Lambda^{-2}) \quad . \end{aligned} \quad (60)$$

The quantity f^2 suggesting a decay constant is actually an operator containing higher derivatives:

$$f^2 = -\frac{N_c \kappa_2 \Lambda^2}{2\pi^2} + \frac{N_c}{12\pi^2} \partial^2 - \frac{N_c \kappa_{-2}}{24\pi^2 \Lambda^2} \partial^2 \partial^2 - \frac{\kappa_{-2}}{24\pi^2 \Lambda^2} \text{tr}_c G_{\nu\rho} G_{\nu\rho} + \dots \quad (61)$$

The dots denote higher-order gluonic terms, derivatives and combinations of them divided by suitable powers of Λ . As such, f^2 summarizes all contributions of second order in A_μ in the chiral Jacobian.

3.1 INDUCED REGULARIZATION

It is interesting to note that not only have we automatically generated, after a trivial rescaling of $\theta(x)$, a standard canonical kinetic energy term for this new field⁴, we actually have a *Pauli-Villars regularized* version of it. This regularization is not put in by hand, but induced by

⁴It seems that the kinetic term has the wrong sign. However, we are dealing with a regularized theory and thus with *bare* parameters; there is no reason why these parameters should have a definite sign. Indeed, the number κ_2 is completely scheme dependent and may have an arbitrary sign.

the chiral rotation, and its associated regularized Jacobian. To see this, let us concentrate on the terms in (60) that are quadratic in θ , and let us furthermore neglect the gluonic terms in the expression (61) for f^2 . Suppose we introduce a pseudoscalar field η_0 with the canonical dimension of *mass* as

$$\theta = \frac{1}{\sqrt{N_f f_0}} \eta_0 \quad (62)$$

and assume that f_0 can be viewed as a bare coupling. Dimensionally, this coupling also goes as $[mass]^1$, and must hence be proportional to the ultraviolet cut-off Λ , the only scale in this regularized theory. Indeed, identifying f_0^2 with the leading term in (61),

$$f_0^2 = -\frac{N_c \kappa_2 \Lambda^2}{2\pi^2} \quad (63)$$

leads to a Lagrangian typical for a pseudoscalar field:

$$\mathcal{L}_{\eta_0} = \frac{1}{2} \partial_\mu \eta_0 \partial_\mu \eta_0 - A_\mu \sqrt{N_f f_0} \partial_\mu \eta_0 + \dots \quad (64)$$

The dots denote higher derivative terms, gluonic terms and self interactions of the pseudoscalar field. Note that f_0 is both cut-off dependent and scheme dependent.

Let us now look at the higher derivative terms. In a perturbative sense, the propagator for the field η_0 can be derived from the bilinear part of $\mathcal{L}'$; keeping also the first of the higher derivative terms of f^2 in (61), it would be, in a momentum representation,

$$G(p^2) = \frac{1}{p^2} - \frac{1}{p^2 + 6\kappa_2 \Lambda^2} . \quad (65)$$

This is a Pauli-Villars regularized bosonic propagator with the regulator mass proportional to Λ^2 , the fermionic cutoff parameter. Actually, in four dimensions we will encounter both quadratic and logarithmic divergences, and one Pauli-Villars regulator therefore does not suffice to regularize the theory completely. This extra Pauli-Villars regulator mass for θ is *also* automatically provided by the chiral rotation. Going to one order higher in the expansion of the free part of f^2 , we find a term which in momentum space is of the form p^4/Λ^4 . The inclusion of this term is equivalent to the addition of one more Pauli-Villars regulator field in the θ -sector.

One may well ask why we stop the Λ^{-1} -expansion at this point, since it seems to lead to such drastic consequences as introducing two Pauli-Villars regulator masses to order $\mathcal{O}(\Lambda^{-2})$. What if we considered yet one more term in the expansion, would that not significantly alter the results? The answer is that the two Pauli-Villars regulators we have included are already sufficient to regularize the ultraviolet divergences associated with the θ -dynamics (at least in perturbation theory). We could of course add yet one higher order if we wished, and also include it in the kinetic energy term instead of treating it as a perturbation. This corresponds to an oversubtracted Pauli-Villars regularization, a valid but inconvenient regularization scheme if we send the cut-off Λ to infinity in the end. Then the θ -propagator with just these two Pauli-Villars regulator terms included is enough to regularize the theory. All higher-order terms are then not just formally, but truly (at least within perturbation theory) suppressed by extra powers of Λ^{-1} . Of course, if we do not wish to send the cut-off Λ to infinity, then all these extra terms in f^2 must be kept for consistency.

The same phenomenon of induced regularization for the collective field occurs in the (1+1)-dimensional case as well [14], and there is probably a simple intuitive reason for it.

The point is that we are throughout performing field transformations within a *regularized* fermionic path integral: all ultraviolet divergences have explicitly been removed. Even after a series of field-enlarging transformations (and the required gauge fixing of the new local symmetry) of such a form that we end up with new propagating fields, the generating functional must still be ultraviolet regularized. The fermionic Jacobian precisely reflects this fact by inducing appropriate regulator terms for the new fields where short-distance singularities threaten to appear.

The above discussion has not covered the gluonic contributions to f^2 in (61). If these were external fields, then we can deduce from the arguments given above that these terms are suppressed by inverse powers of the cut-off. What changes if these are dynamical fields, as in QCD? In this case we know that there may occur gluon condensates like $\langle G^2 \rangle$. In regularized QCD the latter is necessarily of order Λ^4 and will thus contribute to the non-derivative part of f^2 . Simple dimensional reasoning leads to the conclusion that there are similar contributions to the derivative terms in f^2 . In principle, the higher-order gluon condensates have to be included in the same manner. All these terms are calculable from the $1/\Lambda$ -expansion of the chiral Jacobian. We will come back to these gluonic terms in section 3.3.

3.2 AN ANOMALOUS GAUGE FIXING IN FOUR DIMENSIONS

Useful forms of the effective Lagrangian are derived by judicious choices of gauge fixing. Whereas the gauge $\theta(x) = 0$ trivially gives us back cut-off QCD in its original formulation, essentially all other gauge choices that remove part of the QCD degrees of freedom will lead to non-trivial effective Lagrangians. We shall here use some of the lessons learned in the previous 2-dimensional example, and consider the expectation value of the axial singlet current as given by a functional derivative with respect to A_μ at $A_\mu = 0$. It reads

$$i\langle \bar{\psi} \gamma_\mu \gamma_5 \psi \rangle = i\langle \bar{\chi} \gamma_\mu \gamma_5 \chi \rangle + N_f f^2 \partial_\mu \theta + \dots \quad (66)$$

The additional terms represented by dots are at least of third order in $\theta(x)$. The whole expression is, of course, gauge invariant; however, the individual components on the r.h.s. are gauge dependent. Expectation values in eq. (66) are with respect to the fermionic fields only.

In order to keep the discussion of gauge fixing as simple as possible at this point, let us first ignore all $\mathcal{O}(\Lambda^{-2})$ (and higher) corrections in the operator f^2 , which then reduces to f_0^2 as defined in the last subsection. The corrections neglected include non-trivial gluonic contributions, whose significance we shall discuss later.

A comparison with the solvable two-dimensional case [14] is again useful at this point. There, the equivalent of eq.(66) for the same chiral transformation contained only a term linear in θ . By gauge fixing on a suitable linear combination of the resulting two terms comprising the divergence of the physical gauge-invariant axial current, it is possible to find [14] the (presumably unique) gauge-fixing function that switches smoothly between fermionic and bosonic formulations. Since we at least partially wish to achieve something close to “bosonization” of the low-energy QCD Lagrangian, let us first try to choose a gauge-fixing function Φ analogous to the (1+1)-dimensional case:

$$\Phi = \Delta i \frac{\partial_\mu}{N_f f_0^2 \partial^2} \bar{\chi} \gamma_\mu \gamma_5 \chi + (1 - \Delta) \theta . \quad (67)$$

This choice of gauge follows the same philosophy as in (1+1) dimensions: the θ -dependent part of the divergence of the axial current should describe a fraction Δ of the divergence of

the *physical* axial current expressed through the *original* fermionic degrees of freedom $\bar{\psi}$ and ψ .

The above gauge-fixing function is, however, defined in a very formal manner on account of the inverse Laplacian. Furthermore, if we wish to implement Φ as a delta-function constraint in the path integral, we should be careful that it satisfies correct transformation properties. Under a global chiral rotation,

$$\bar{\chi}(x) \rightarrow \bar{\chi}(x)e^{i\alpha\gamma_5} \quad , \quad \chi(x) \rightarrow e^{i\alpha\gamma_5}\chi(x) \quad , \quad (68)$$

the object involving $\bar{\chi}\gamma_\mu\gamma_5\chi$ remains *classically* invariant but quantum mechanically it shifts due to the chiral anomaly. We find the shift from the expansion given in eq. (60):

$$i\frac{\partial_\mu}{N_f f_0^2 \partial^2} \bar{\chi}\gamma_\mu\gamma_5\chi \rightarrow i\frac{\partial_\mu}{N_f f_0^2 \partial^2} \bar{\chi}\gamma_\mu\gamma_5\chi + \alpha \quad . \quad (69)$$

The formal operator $N_f f_0^2 \partial^2$ is included precisely in order to cancel the additional factors multiplying α .

The action does not remain invariant under constant chiral rotations, but shifts due to the axial anomaly:

$$S' = \int d^4x \mathcal{L}' \rightarrow S' - 2iN_f\alpha \int d^4x \frac{1}{32\pi^2} \epsilon_{\mu\nu\rho\sigma} \text{tr}_c G_{\mu\nu} G_{\rho\sigma} \quad . \quad (70)$$

Assuming that we sum only over integer winding numbers, the action does, however, remain invariant under constant chiral rotations of the form $\alpha = n\pi/N_f$. This means that $\theta(x)$ is only globally defined modulo π/N_f .

Gauge-fixing the constant $\theta(x)$ modes is then again non-trivial. The gauge-fixing constraint must respect the above periodicity property; there must, even in the gauge-fixed path integral, be no distinction between $\theta(x)$ and $\theta(x) + n\pi/N_f$. If we choose a δ -function constraint to implement the gauge choice, this δ -function must then necessarily be *globally periodic*, quite analogous to the 2-dimensional case.

We can again represent such a globally periodic δ -function by means of a Nakanishi-Lautrup auxiliary field $b(x)$, so that this gauge-fixing function just provides a few new terms in the action. However, these new terms in the action will in general modify the relevant chiral Jacobian, and for consistency – or, equivalently, if we wish to keep BRST symmetry in the gauge fixing – new terms must be added to the action to compensate for the change in the Jacobian. This leads to a rather involved procedure of gauge fixing, and we have preferred to present a simpler derivation. The final result agrees with that obtained through the direct method.

This is based on the fact that quantum mechanically we can consider the above gauge-fixing function as derived from a constraint in the *original* representation of QCD of the form

$$\Phi' = \Delta i \frac{\partial_\mu}{N_f f_0^2 \partial^2} \bar{\psi}\gamma_\mu\gamma_5\psi + \theta \quad . \quad (71)$$

We shall here illustrate the gauge-fixing procedure by this shortcut. The equivalence of Φ and Φ' is valid at the level of expectation values. In the final gauge-fixed action there is of course an enormous amount of freedom left, since we can add any term proportional to only $b(x)$ (and higher powers thereof), as long as these terms do not multiply any functions of θ . These terms do not affect physical gauge-invariant Green functions. We shall make use of this freedom shortly.

As mentioned, it is convenient to implement the periodic δ -function by a functional Fourier representation. If we introduce the δ -function constraint in the shortcut manner described above, this corresponds to

$$\begin{aligned}\delta(\Phi') &= \int \mathcal{D}[b] \exp \left[- \int d^4x d^4z \, b(x) \left(\frac{i\Delta}{N_f f_0^2} \partial_{(x-z)}^{-2} \partial_\mu^{(z)} J_\mu^5(z) - \delta(x-z) \theta(z) \right) \right] \\ &= \int \mathcal{D}[b] \exp \left[- \int d^4x \left(\frac{i\Delta}{N_f f_0^2} B_\mu(x) J_\mu^5(x) + \theta(x) \partial_\mu B_\mu(x) \right) \right] ,\end{aligned}\quad (72)$$

where the axial vector field $B_\mu(x)$ is defined by

$$B_\mu(x) \equiv \int d^4y d^4z \, b(y) \partial_{(y-z)}^{-2} \partial_\mu^{(z)} \delta(z-x) . \quad (73)$$

Note that this implies $b(x) = -\partial_\mu B_\mu(x)$ as in the 2-dimensional case. Global periodicity of the δ -function means that b is constrained,

$$\int d^4x \, b(x) = - \int d^4x \, \partial_\mu B_\mu = ik N_f , \quad (74)$$

where k is an arbitrary integer. This global constraint means that $b(x)$ (or $\partial_\mu B_\mu$) share certain properties with topologically non-trivial fields. But we of course have no direct interpretation in terms of gauge potentials. In two dimensions, the same global constraint has a more physical interpretation in terms of an Abelian gauge field of non-trivial topology [14]. Nevertheless, these topological properties of $b(x)$ play a similarly profound rôle in chiral Ward identities of a truncated QCD (to be defined in a precise manner later) as in the 2-dimensional case.

With this implementation of the gauge fixing in the original Lagrangian, we can now perform the same chiral transformation as before. The only change is that in the transformed Lagrangian $\mathcal{L}'$, the external axial vector source A_μ has to be replaced by $A_\mu - \frac{\Delta}{N_f f_0^2} B_\mu$ and a term $\theta \partial_\mu B_\mu$ has to be added.

After gauge-fixing the only remnant of the $U(1)$ axial gauge symmetry is the BRST symmetry

$$\begin{aligned}\delta \bar{\chi}(x) &= i \bar{\chi}(x) \gamma_5 c(x) \\ \delta \chi(x) &= -i c(x) \gamma_5 \chi(x) \\ \delta \theta(x) &= -c(x) \\ \delta c(x) &= 0 \\ \delta \bar{c}(x) &= b(x) \\ \delta b(x) &= 0.\end{aligned}\quad (75)$$

The ghost term is trivial in the present case:

$$\mathcal{L}_{ghost} = \bar{c} c \quad (76)$$

It decouples and will be neglected in the following.

The complete gauge-fixed Lagrangian now reads

$$\mathcal{L}'' = \bar{\chi} \left(\not{\partial} - i \not{A} - \not{M} - i \left(A - \frac{\Delta}{N_f f_0^2} B - \not{\partial} \theta \right) \gamma_5 \right) \chi + \bar{c} c + \mathcal{L}_{YM}$$

$$\begin{aligned}
& + \frac{N_f f_0^2}{2} \partial_\mu \theta \partial_\mu \theta - N_f f_0^2 A_\mu \partial_\mu \theta - (1 - \Delta) \theta \partial_\mu B_\mu \\
& - \frac{N_f N_c}{12\pi^2} \left(\left((A_\mu - \frac{\Delta}{N_f f_0^2} B_\mu - \partial_\mu \theta) (A_\mu - \frac{\Delta}{N_f f_0^2} B_\mu - \partial_\mu \theta) \right)^2 \right. \\
& \left. - \left((A_\mu - \frac{\Delta}{N_f f_0^2} B_\mu) (A_\mu - \frac{\Delta}{N_f f_0^2} B_\mu) \right)^2 \right) \\
& - \theta \frac{i N_f}{16\pi^2} \epsilon_{\mu\nu\rho\sigma} \left(\text{tr}_c G_{\mu\nu} G_{\rho\sigma} - 4N_c \partial_\mu V_\nu \partial_\rho V_\sigma - \frac{4N_c}{3} \partial_\mu A_\nu \partial_\rho A_\sigma \right) \\
& + \mathcal{O}(\Lambda^{-2}) .
\end{aligned} \tag{77}$$

Due to its longitudinal character B_μ does not appear in the WZ-like term of the last line.

It is convenient to choose a slightly different gauge which will only affect higher order correlation functions of our gauge fixing expression Φ' . As already mentioned we are allowed to add arbitrary terms containing $b(x)$ or powers thereof as long as they are not coupled to $\theta(x)$. To be precise, we choose them such that the gauge fixed Lagrangian looks like

$$\begin{aligned}
\mathcal{L}'' &= \bar{\chi} \left(\not{\partial} - i\not{A} - i\not{M} - i \left(\not{A} - \frac{\Delta}{N_f f_0^2} \not{B} - \not{\partial} \theta \right) \gamma_5 \right) \chi + \bar{c}c + \mathcal{L}_{YM} \\
&+ \frac{N_f f_0^2}{2} \partial_\mu \theta \partial_\mu \theta - N_f f_0^2 A_\mu \partial_\mu \theta - (1 - \Delta) \theta \partial_\mu B_\mu \\
&- \frac{N_f N_c}{12\pi^2} \left(\left((A_\mu - \frac{\Delta}{N_f f_0^2} B_\mu - \partial_\mu \theta) (A_\mu - \frac{\Delta}{N_f f_0^2} B_\mu - \partial_\mu \theta) \right)^2 - (A_\mu A_\mu)^2 \right) \\
&- \theta \frac{i N_f}{16\pi^2} \epsilon_{\mu\nu\rho\sigma} \left(\text{tr}_c G_{\mu\nu} G_{\rho\sigma} - 4N_c \partial_\mu V_\nu \partial_\rho V_\sigma - \frac{4N_c}{3} \partial_\mu A_\nu \partial_\rho A_\sigma \right) \\
&+ \mathcal{O}(\Lambda^{-2})
\end{aligned} \tag{78}$$

3.3 THE $\Delta = 1$ GAUGE

In two space-time dimensions, the case $\Delta = 1$ corresponds to the “bosonization gauge” [14]. It is therefore natural to consider the corresponding analogue here in four dimensions. Of course, we have no hope of completely bosonizing QCD, but we should not be surprised to find that the corresponding $\Delta = 1$ gauge in four dimensions most conveniently extracts the pseudoscalar collective degrees of freedom. Actually, we can only bosonize those degrees of freedom representing the longitudinal component of $J_{5\mu}$. With $\Delta = 1$, $\mathcal{L}''$ becomes

$$\begin{aligned}
\mathcal{L}'' &= \bar{\chi} \left(\not{\partial} - i\not{A} - i\not{M} - i \left(\not{A} - \frac{1}{N_f f_0^2} \not{B} - \not{\partial} \theta \right) \gamma_5 \right) \chi + \bar{c}c + \mathcal{L}_{YM} \\
&+ \frac{N_f f_0^2}{2} \partial_\mu \theta \partial_\mu \theta - N_f f_0^2 A_\mu \partial_\mu \theta \\
&- \frac{N_f N_c}{12\pi^2} \left(\left((A_\mu - \frac{1}{N_f f_0^2} B_\mu - \partial_\mu \theta) (A_\mu - \frac{1}{N_f f_0^2} B_\mu - \partial_\mu \theta) \right)^2 - (A_\mu A_\mu)^2 \right) \\
&- \theta \frac{i N_f}{16\pi^2} \epsilon_{\mu\nu\rho\sigma} \left(\text{tr}_c G_{\mu\nu} G_{\rho\sigma} - 4N_c \partial_\mu V_\nu \partial_\rho V_\sigma - \frac{4N_c}{3} \partial_\mu A_\nu \partial_\rho A_\sigma \right) \\
&+ \mathcal{O}(\Lambda^{-2}) .
\end{aligned} \tag{79}$$

As expected, this involves all fields – fermionic and bosonic – interacting in all respects. We are nowhere near a bosonized form of the QCD Lagrangian. But, as we shall see, many of the simplifying features of two-dimensional bosonization nevertheless remain hidden in this form of the Lagrangian.

In order to view (79) as an effective Lagrangian, we need additional input. The obvious choice would be to identify the θ -field with the flavour-singlet pseudoscalar field of the η' meson, in appropriate units. But depending on the questions we wish to answer, such an explicit identification may not be necessary. Certainly, eq. (79) gives in the $\Lambda \rightarrow \infty$ limit the correct QCD action for describing the dynamics of the composite operator $J_\mu^5(x) = i\bar{\psi}\gamma_\mu\gamma_5\psi(x)$ of the original quark fields supposing the limit for the gluonic terms of the chiral Jacobian is taken in an appropriate way. Taking one partial derivative, we can equally well describe $\partial_\mu J_\mu^5(x)$, which is a non-zero operator due to the chiral anomaly. It has quantum numbers $J^{PC} = 0^{-+}$, and is a singlet under flavour. As such, this object should have a non-vanishing overlap with the physical η' meson. For example, if we were able to compute the long-distance fall-off of the corresponding two-point correlation function, this should provide us with the mass of the lowest-lying state of these quantum numbers. Ignoring the possibility of a lighter glueball with the same quantum numbers, this is the mass of the η' meson.

Let us now proceed with such a calculation. Going back to the defining equation (51), we note that the connected 2-point function of $\partial_\mu J_\mu^5(x)$ can be obtained by differentiating twice with respect to a pseudoscalar source $\sigma(x)$ defined by splitting $A_\mu = \partial_\mu\sigma(x) + A_\mu^T$ into a longitudinal and a transverse part. Shifting B_μ

$$B_\mu(x) \rightarrow B_\mu(x) + N_f f_0^2 \partial_\mu \theta(x) - N_f f_0^2 \partial_\mu \sigma(x) \quad (80)$$

leads to a Lagrangian

$$\begin{aligned} \mathcal{L}''' = & \bar{\chi} \left(\not{\partial} - i\not{G} - i \left(A^T - \frac{1}{N_f f_0^2} B \right) \gamma_5 \right) \chi + \mathcal{L}_{ghost} + \mathcal{L}_{YM} \\ & - \frac{N_f N_c}{12\pi^2} \left(\left((A_\mu^T - \frac{1}{N_f f_0^2} B_\mu)(A_\mu^T - \frac{1}{N_f f_0^2} B_\mu) \right)^2 - (A_\mu A_\mu)^2 \right) \\ & + \frac{N_f f_0^2}{2} \partial_\mu \theta \partial_\mu \theta - N_f f_0^2 \partial_\mu \sigma \partial_\mu \theta \\ & - \theta \frac{i N_f}{16\pi^2} \epsilon_{\mu\nu\rho\sigma} \left(\text{tr}_c G_{\mu\nu} G_{\rho\sigma} - 4N_c \partial_\mu V_\nu \partial_\rho V_\sigma - \frac{4N_c}{3} \partial_\mu A_\nu^T \partial_\rho A_\sigma^T \right) \\ & + \mathcal{O}(\Lambda^{-2}) \end{aligned} \quad (81)$$

Apart from contact terms only a linear coupling of σ to θ is left. The remaining part of $\mathcal{O}(\Lambda^{-2})$ is also independent of θ because it contains θ only in the combination $B_\mu + N_f f_0^2 \partial_\mu \theta$; after the shift (80) θ disappears from these terms.

We can now derive some exact Ward identities, setting the external sources to zero: The original anomalous Ward identity

$$\partial_\mu \langle J_{5\mu} \rangle = i \partial_\mu \langle \bar{\psi} \gamma_\mu \gamma_5 \psi \rangle = -i \frac{N_f}{16\pi^2} \langle G\tilde{G} \rangle + \mathcal{O}(\Lambda^{-2}) \quad (82)$$

is now just the equation of motion for the field θ :

$$f_0^2 \partial^2 \langle \theta \rangle = -i \frac{1}{16\pi^2} \langle G\tilde{G} \rangle + \mathcal{O}(\Lambda^{-2}) \quad (83)$$

with the shorthand notation

$$G\tilde{G}(x) = \epsilon_{\mu\nu\rho\sigma} \text{tr}_c G_{\mu\nu}(x) G_{\rho\sigma}(x) \quad (84)$$

Analogously we find an anomalous expression for the 2-point function in the original QCD representation,

$$\begin{aligned} \langle \partial_\mu J_{5\mu}(x) \partial_\nu J_{5\nu}(y) \rangle &= - \left(\frac{N_f}{16\pi^2} \right)^2 \langle G\tilde{G}(x) G\tilde{G}(y) \rangle \\ &\quad - N_f f_0^2 \partial^2 \delta(x-y) + \mathcal{O}(\Lambda^{-2}) . \end{aligned} \quad (85)$$

Note the presence of a contact term in eq. (85). We will come back to it at the end of this subsection. The same identity can be derived considering a simple infinitesimal shift of θ to second order:

$$\begin{aligned} f_0^4 \langle \partial^2 \theta(x) \partial^2 \theta(y) \rangle &= - \left(\frac{1}{16\pi^2} \right)^2 \langle G\tilde{G}(x) G\tilde{G}(y) \rangle \\ &\quad - f_0^2 \partial^2 \delta(x-y) + \mathcal{O}(\Lambda^{-2}) \end{aligned} \quad (86)$$

These two examples illustrate that for appropriate Green functions our $\Delta = 1$ gauge really identifies

$$\partial_\mu J_{5\mu} \sim N_f f_0^2 \partial^2 \theta \quad (87)$$

as *operators*. Actually, this identification holds only up to fourth order correlation functions. The reason is that – as already mentioned – we have modified our original gauge by the inclusion of terms depending on $b(x)$ (resp. $B_\mu(x)$). These terms were precisely of fourth and higher order.

The gauge-fixing procedure presented above can therefore be understood from another point of view. Essentially it amounts to introducing explicitly, at the Lagrangian level, an “interpolating” field according to the relation

$$\partial_\mu J_{5\mu} = \sqrt{N_f} f_0 \partial^2 \eta_0 , \quad (88)$$

where f_0 is the corresponding decay constant. On mass shell, $p^2 = m_{\eta_0}^2$, this would indeed be a conventional definition of an interpolating field. In our gauge, however, the interpolating field is identified even off-shell, as it should be in a path integral framework.

Note that we are not claiming, and need not claim, that the field $\eta_0(x)$ is to be identified with the η' meson. Only at large distances, where other isosinglet pseudoscalar states are suppressed because of their presumed larger mass, can we indirectly make such an identification, as in the usual reduction formalism. The present gauge choice simply enforces that a portion Δ (and in this particular case of $\Delta = 1$, *all*) of the divergence of the physical axial current $J_{5\mu} = i\bar{\psi}\gamma_\mu\gamma_5\psi$ is carried by the collective field.

We now formally integrate out all fields in (81) except θ to arrive at an effective Lagrangian

$$\mathcal{L}_{eff} = \frac{F_0^2}{2} \partial_\mu \theta \partial_\mu \theta + \frac{F_0^2 M_0^2}{2} \theta^2 + \dots \quad (89)$$

The dots denote higher derivative terms and self-interactions of order θ^3 . The parameters F_0 and M_0 are defined through

$$F_0^2 M_0^2 = \int d^4x \left\langle \frac{N_f}{16\pi^2} G\tilde{G}(x) \frac{N_f}{16\pi^2} G\tilde{G}(0) \right\rangle_{trunc} \quad (90)$$

and

$$F_0^2 = N_f f_0^2 - \int d^4x \frac{x^2}{8} \left\langle \frac{N_f}{16\pi^2} G\tilde{G}(x) \frac{N_f}{16\pi^2} G\tilde{G}(0) \right\rangle_{trunc} . \quad (91)$$

The parameters F_0, M_0 have to be interpreted as *bare* ones. The expression for $F_0^2 M_0^2$ in (90) is formally the same as derived by Witten [16] and Veneziano [17] in the limit $N_c \rightarrow \infty$. In our case this relation is valid for arbitrary N_c . But the expectation values $\langle \dots \rangle_{trunc}$ have to be taken with respect to a “truncated” version of QCD:

$$\begin{aligned} \mathcal{L}_{trunc} = & \bar{\chi} \left(\not{\partial} - i\not{A} + i \frac{1}{N_f f_0^2} \not{B} \gamma_5 \right) \chi + \mathcal{L}_{ghost} + \mathcal{L}_{YM} \\ & - \frac{N_c}{12\pi^2 N_f^3 f_0^8} (B_\mu B_\mu)^2 \\ & + \mathcal{O}(\Lambda^{-2}) , \end{aligned} \quad (92)$$

where we have neglected external sources.

At first glance one would argue that the topological susceptibility has to be zero in such a theory because of the massless quarks. If this were true, then our field θ would be massless. Indeed, for the full theory we can derive from the Ward identities (86) that

$$\int d^4x \langle G\tilde{G}(x) G\tilde{G}(0) \rangle = 0 . \quad (93)$$

This is a well-known result in massless QCD. Let us do an analogous step for the truncated theory. In particular we get from a relation resembling (86) the identity

$$\begin{aligned} \int d^4x \langle \partial_\mu B_\mu(x) \partial_\nu B_\nu(0) \rangle_{trunc} = & - \left(\frac{N_f}{16\pi^2} \right)^2 \int d^4x \langle G\tilde{G}(x) G\tilde{G}(0) \rangle_{trunc} \\ & + \mathcal{O}(\Lambda^{-2}) , \end{aligned} \quad (94)$$

the analogue of relation (50) in two dimensions. Now we see the importance of gauge-fixing the zero-modes in a proper manner: it forced us to introduce the non-trivial constraint $\int d^4x \partial_\mu B_\mu(x) = const..$ It is precisely this constraint that may imply a non-vanishing topological susceptibility in the truncated theory and thus a nonvanishing mass for the field θ . Without this constraint, one could decouple the B_μ -field from the fermions through a chiral rotation. The standard proof of a vanishing topological susceptibility in the theory with massless quarks would then go through. However, when the B_μ -field is constrained in the manner shown, this decoupling by means of a chiral rotation is impossible [29], and the topological susceptibility may be non-vanishing even (as in this truncated theory) when the fermions are massless. The field B_μ is probably the closest one can get at an explicit QCD Lagrangian realization of the Veneziano ghost [17].

The essence of this exercise was to get some insight into the details of the gauge-fixing mechanism. Let us now try to understand these results from a more phenomenological point of view. For that purpose it is useful to compare with the argumentation of Witten in ref.[16], where it was argued that the vanishing of the topological susceptibility in massless QCD can be explained as a cancellation of the pure gluonic part by a contribution from the η' meson. In our case, the appearance of a non-zero topological susceptibility in the *truncated* QCD would be interpreted as an effect of the b -field interaction. This additional interaction, which arises through the gauge constraint, can be thought of as removing the η' -part of the topological susceptibility. The picture of the mass generation of the η' for finite N_c that emerges then seems to coincide with the one of ref. [25]. In fact, we see here one explicit Lagrangian realization of what is meant by the notion of “no η' insertion”, which plays such a crucial rôle in ref.[25]. In our formulation this subtraction of the η' degree of freedom from the QCD generating functional is done in an exact manner; the expectation values with no

η' insertions simply mean expectation values taken with respect to the *chirally rotated* quark fields $\bar{\chi}, \chi$ and the gluons.

At the moment, we have not specified how to renormalize the effective theory. A major difficulty with the present approach is that everything is expressed in terms of bare parameters in the cut-off theory. Being explicitly cut-off dependent, we are not surprised to find that the coefficients of the effective couplings are also scheme dependent. The whole set of effective one-loop interactions between the bosonic collective fields and left-over QCD degrees of freedom indeed follow directly from the Pauli-Villars regulator fields. This is just as in the solvable case of two dimensions [14].

In principle, the renormalization prescription should be equivalent to the one for QCD in its original representation. The situation is, however, further complicated by the fact that we are necessarily dealing with a generating functional of composite operators. One renormalization procedure including arbitrary insertions of such operators has recently been suggested in ref.[26] (see also ref. [27]). As it stands, the unrenormalized theory has, with massless quarks, only one mass scale: that of the cut-off Λ . This means that all dimensionful couplings in the effective theory are given by powers of this ultraviolet cut-off. In the renormalized theory this cut-off becomes replaced by a physical mass scale, to be extracted from experiments. In the end, if one integrates out all gluonic and quark degrees of freedom and leaves only the collective fields, the physical couplings are directly related to gluonic and fermionic correlators, moments thereof, and condensates. The precise relationship between the couplings of the collective field Lagrangian and these vacuum expectation values will be of roughly the kind discussed in the case of the Witten-Veneziano relation above, but will of course require a non-perturbative treatment. The fact that physical couplings will be related to these Green functions is also evident if we return to the definition of the gauge-fixing condition (67) in subsection 3.2.. The term containing f_0^2 should in fact depend on f^2 with contributions also from gluonic fields, and a full treatment should incorporate the effect of integrating out the gluonic degrees of freedom. Intuitively, we would expect that one major effect of such a renormalization program would relate f^2 to gluonic condensates.

An important point needs to be mentioned here. If we turn off all gluonic interactions, we can still go through the steps of deriving the “effective Lagrangian” by means of chiral rotations in the cut-off theory. It appears as if the only difference between these two effective Lagrangians arises from “small gluonic corrections”. Ignoring complications due to the anomalous gauge fixing, this would seem to imply that a convenient effective Lagrangian for a theory of *free* fermions is of essentially the same form as the one derived here for QCD! The same argument can, incidentally, be raised against the purely bosonic effective Lagrangians of refs. [9, 10]. What is the resolution of this apparent contradiction? As we have emphasized earlier, the collective field technique is not tied to such notions as spontaneous symmetry breaking and the existence of (pseudo-) Goldstone bosons. We can *always* introduce given collective fields in a theory, independently of whether these collective fields may be related to asymptotic states of the theory. For a free theory, a collective field representation is valid, but of limited use: The corresponding collective field mode will decay into the free constituents. Alternatively, we can look at this from the point of view of renormalized couplings in the effective Lagrangian. For a free theory, the only scale with which to specify effective couplings will remain the (unphysical) ultraviolet cut-off. In the full theory of QCD, these effective couplings will be specified by dynamics, including such crucial features as chiral symmetry breaking, gluonic condensates etc. What this shows very explicitly is that the term “gluonic corrections” may be quite misleading; in the end a large

fraction of the effective couplings of the resulting chiral Lagrangian for the collective field may be given by the values of gluonic condensates, moments of such condensates, the chiral condensate, and so on.

Note that in the limit $\Lambda \rightarrow \infty$, the generating functional with the Lagrangian in eq. (79) is an *exact* rewrite of the generating functional of QCD, eq. (51). No approximations are involved at this point as long as we perform the limit $\Lambda \rightarrow \infty$ for the whole expansion of the chiral Jacobian in a careful manner, taking into account possible gluonic condensates. Even for finite Λ can we, in principle, make the representation as accurate as we wish by including a sufficient number of terms in the known $1/\Lambda$ expansion. In this sense, the rewrite of the generating functional is always exact, for any value of Λ .

We emphasize that a non-perturbative derivation of the physical couplings is possible in principle, since the cut-off effective Lagrangian contains the same information as the cut-off QCD Lagrangian in terms of the original quark and gluon degrees of freedom. As in the case of QCD, also the effective Lagrangian is amenable to numerical studies from which the effective couplings can be extracted.

4 Constituent Quarks and Mesonic Degrees of Freedom

In this section we shall briefly outline the generalization of the present effective Lagrangian technique to the case of the $SU(N_f)$ pseudoscalar multiplet. In contrast with the $U(1)$ case discussed above, the flavour non-singlet axial currents are exactly conserved in the limit of massless quarks. We need not be concerned with the question of spontaneous chiral symmetry breaking here; the effects will automatically be taken into account.

How do we introduce the appropriate collective fields for this non-Abelian (flavoured) case? As before, the main input is the choice of quantum numbers we wish to describe. We then start again with a generating functional of QCD,

$$\begin{aligned}\mathcal{Z}[V, A] &= \int \mathcal{D}[\bar{q}, q] d\mu[G] e^{-\int d^4x \mathcal{L}} \\ \mathcal{L} &= \bar{q}(\not{\partial} - i\not{G} - i\not{V} - i\not{A}\gamma_5)q + \mathcal{L}_{YM},\end{aligned}\tag{95}$$

where we now consider external sources V_μ and A_μ that are elements of $SU(N_f)$.

For technical reasons that will become clear shortly, it is most convenient to introduce collective fields $\theta(x)$ by, *e.g.*, purely left-handed transformations:

$$q_L(x) = e^{2i\theta(x)}\chi_L(x) \quad , \quad \bar{q}_L(x) = \bar{\chi}_L(x)e^{-2i\theta(x)}\tag{96}$$

i.e. local phase transformations acting only on the left-handed spinors:

$$q_L = P_+ q \quad , \quad \bar{q}_L = \bar{q} P_- \quad , \quad P_\pm = \frac{1}{2}(1 \pm \gamma_5) .\tag{97}$$

The transformation (96) is a combination of an axial and a pure vector gauge transformation. The latter can easily be corrected for by a suitable opposite transformation; there are no complications due to regularization, because we are using a scheme that preserves vector gauge invariance.

Here $\theta(x)$ is understood to be an element of a Lie algebra, in particular that of $SU(N_f)$,

$$\theta = \theta^a \lambda^a\tag{98}$$

where λ^a are the generators of $SU(N_f)$ with the convention $\text{tr } \lambda^a \lambda^b = \delta^{ab}$. It is convenient to use the combinations

$$L_\mu = V_\mu + A_\mu \quad , \quad R_\mu = V_\mu - A_\mu \quad (99)$$

instead of V_μ and A_μ .

The transformation (96) exhibits a change of the regularized fermionic functional integral measure due to its handedness. In order to calculate the corresponding contribution to the Lagrangian we again use a Pauli-Villars regularization for reasons described in section 2. The generating functional now reads

$$\begin{aligned} \mathcal{Z}_\Lambda[V, A] &= \int \mathcal{D}_\Lambda[\bar{\chi}, \chi] d\mu[G] e^{-\int d^4x \mathcal{L}'} \\ \mathcal{L}' &= \bar{\chi} \gamma_\mu (\partial_\mu - iG_\mu - iL_\mu^\theta P_+ - iR_\mu P_-) \chi + \mathcal{L}_J + \mathcal{L}_{WZ} + \mathcal{L}_{YM} , \end{aligned} \quad (100)$$

where only L_μ is modified as

$$L_\mu^\theta = U^\dagger L_\mu U + iU^\dagger \partial_\mu U . \quad (101)$$

We have now introduced the common symbol U as

$$U(x) = e^{2i\theta(x)} \quad U^\dagger(x) = e^{-2i\theta(x)} . \quad (102)$$

Again, $\mathcal{L}_J$ and $\mathcal{L}_{WZ}$ are induced by the fermionic measure.

The positive parity part can, as in the abelian case, be ordered as an expansion in inverse powers of the ultraviolet cut-off Λ :

$$\mathcal{L}_J = \Lambda^2 \mathcal{L}_2 + \mathcal{L}_0 + \frac{1}{\Lambda^2} \mathcal{L}_{-2} + \frac{1}{\Lambda^4} \mathcal{L}_{-4} + \dots \quad (103)$$

where the first three terms are given by

$$\begin{aligned} \mathcal{L}_2 &= \frac{N_c \kappa_2}{4\pi^2} \text{tr}_f A_\mu^{(s)} A_\mu^{(s)} \Big|_{s=1}^0 \\ \mathcal{L}_0 &= \frac{N_c}{8\pi^2} \text{tr}_f \left(-iF_{\mu\nu}^{(s)} [A_\mu^{(s)}, A_\nu^{(s)}] + \frac{1}{3} D_\mu^{(s)} A_\nu^{(s)} D_\mu^{(s)} A_\nu^{(s)} - \frac{2}{3} (A_\mu^{(s)} A_\mu^{(s)})^2 \right. \\ &\quad \left. + \frac{4}{3} A_\mu^{(s)} A_\nu^{(s)} A_\mu^{(s)} A_\nu^{(s)} \right) \Big|_{s=1}^0 \\ \mathcal{L}_{-2} &= \frac{\kappa_{-2}}{48\pi^2} \text{tr}_f \left(N_c \partial^2 A_\mu^{(s)} \partial^2 A_\mu^{(s)} + A_\mu^{(s)} A_\mu^{(s)} \text{tr}_c G_{\nu\rho} G_{\nu\rho} + \dots \right) \Big|_{s=1}^0 . \end{aligned} \quad (104)$$

Here, tr_f and tr_c denote the traces over flavour and colour indices. The terms omitted in $\mathcal{L}_{-2}$ and denoted by dots are at least fourth order in $A_\mu^{(s)}$ and $V_\mu^{(s)}$. The expression for $\mathcal{L}_{-4}$ is quite lengthy, and we do not reproduce it here.

The leading term of the negative parity part is the integrated Bardeen-anomaly:

$$\begin{aligned} \mathcal{L}_{WZ} &= \frac{i}{16\pi^2} \int_1^0 ds \epsilon_{\mu\nu\rho\sigma} \text{tr}_f \text{tr}_c \theta \left(F_{\mu\nu}^{(s)} F_{\rho\sigma}^{(s)} + \frac{1}{3} A_{\mu\nu}^{(s)} A_{\rho\sigma}^{(s)} \right. \\ &\quad \left. + \frac{8i}{3} (F_{\mu\nu}^{(s)} A_\rho^{(s)} A_\sigma^{(s)} + A_\mu^{(s)} F_{\nu\rho}^{(s)} A_\sigma^{(s)} + A_\mu^{(s)} A_\nu^{(s)} F_{\rho\sigma}^{(s)}) \right. \\ &\quad \left. + \frac{32}{3} A_\mu^{(s)} A_\nu^{(s)} A_\rho^{(s)} A_\sigma^{(s)} \right) + \mathcal{O}(\Lambda^{-2}) . \end{aligned} \quad (105)$$

The covariant derivatives and field strength tensors appearing in eqs. (104) and (105) are defined as

$$\mathcal{D}_\mu A_\nu = \partial_\mu A_\nu - i[V_\mu, A_\nu]$$

$$\begin{aligned} A_{\mu\nu} &= \mathcal{D}_{[\mu} A_{\nu]} \\ F_{\mu\nu} &= \partial_{[\mu} V_{\nu]} - i[V_{\mu}, V_{\nu}] - i[A_{\mu}, A_{\nu}] \end{aligned} \quad (106)$$

and the transformed fields appearing in (104) and (105) as

$$\begin{aligned} V_{\mu}^{(s)} &= \frac{1}{2} \left(e^{-2is\theta} L_{\mu} e^{2is\theta} + i e^{-2is\theta} \partial_{\mu} e^{2is\theta} + R_{\mu} \right) \\ A_{\mu}^{(s)} &= \frac{1}{2} \left(e^{-2is\theta} L_{\mu} e^{2is\theta} + i e^{-2is\theta} \partial_{\mu} e^{2is\theta} - R_{\mu} \right) . \end{aligned} \quad (107)$$

The parameter s ranging from 0 to 1 thus defines a continuous transformation which, for $s = 1$, coincides with (96).

If we now declare $\theta(x)$ a dynamical variable by integrating over the invariant Haar measure $\int \mathcal{D}[U]$, a new local non-Abelian gauge symmetry emerges:

$$\begin{aligned} \chi_L(x) &\rightarrow e^{2i\alpha(x)} \chi_L(x) \\ \bar{\chi}_L(x) &\rightarrow \bar{\chi}_L(x) e^{-2i\alpha(x)} \\ U(x) &\rightarrow U(x) e^{-2i\alpha(x)} \end{aligned} \quad (108)$$

where $\alpha(x)$ is now a (local) transformation parameter in the same representation of $SU(N_f)$ as $\theta(x)$.

As in the Abelian case, we discover an induced regularization, actually due to the same operator f^2 as in (61). This can be seen by extracting from $\mathcal{L}_J$ the leading quadratic term in $U^\dagger \partial_{\mu} U$. In particular, it reads

$$\mathcal{L}_J = -\frac{1}{8} \text{tr} U^\dagger \partial_{\mu} U f^2 U^\dagger \partial_{\mu} U + \dots \quad (109)$$

The omitted terms are of higher order in $U^\dagger \partial_{\mu} U$ or contain external sources.

As in the flavour-singlet case, the crucial step now is the choice of gauge fixing. To derive a convenient form of the effective Lagrangian, the optimal solution would be to find a suitable operator $\mathcal{O}_i$ in terms of the original quark fields $\bar{\psi}$ and ψ such that the quantum version of the same operator expressed in terms of the chirally rotated quark fields $\bar{\chi}, \chi$ and the correction due to the chiral field U separates. Then one can choose the very convenient gauge in which $\mathcal{O}_i[\bar{\chi}, \chi] = 0$. In the flavour-singlet case, the chiral Jacobian precisely allows us to find such a gauge; in that case there is only a gauge fixing at the quantum level, and the above separation of variables is guaranteed. Analogues of such a gauge fixing can be found in the flavoured case too. Without carrying such a gauge fixing through at this point, what will be the rough form of the resulting gauge-fixed Lagrangian? It is interesting to compare it with what has become known as the constituent chiral quark model of Manohar and Georgi [7] (see also the last part of ref. [5]).

In its original formulation [7] and with our conventions, the chiral quark model is described in terms of a Lagrangian

$$\mathcal{L} = \bar{\psi}(i\not{D} + \not{p} + g_A \not{p} \gamma_5 - m)\psi + \frac{1}{4g^2} \text{tr} G_{\mu\nu} G_{\mu\nu} + \frac{1}{8} f^2 \text{tr} \partial_{\mu} U^\dagger \partial_{\mu} U + \dots \quad (110)$$

Here, a_{μ} and v_{μ} are given as

$$a_{\mu} = \frac{i}{2} (\xi^\dagger \partial_{\mu} \xi - \xi \partial_{\mu} \xi^\dagger) , \quad v_{\mu} = \frac{i}{2} (\xi^\dagger \partial_{\mu} \xi + \xi \partial_{\mu} \xi^\dagger) \quad (111)$$

and $U(x) = \xi(x)\xi(x)$. Up to an ordinary gauge transformation with ξ the expressions for a_μ, v_μ coincide with those in (107) for $s = 1$. External sources have not been included. The covariant derivative D_μ is just the usual $D_\mu = \partial_\mu - iG_\mu$ of QCD. In the manner written above, this appears to be the standard QCD Lagrangian, with the extra chiral-model terms (and as yet undetermined chiral coupling g_A) added artificially on top. The reason why this is argued not to overcount the degrees of freedom [7] is that the quarks are to be viewed as “constituent” quarks in eq. (110). For example, the quark mass term above is not the mass term of the current quarks, but rather a number on the order of 350 MeV.

It would be tempting to identify our rotated and gauge-fixed fermion fields $\bar{\chi}$ and χ with the constituent quarks of the Lagrangian (110). Before doing this, we should, however, recall that the fields $\bar{\chi}, \chi$ have not been introduced for the purpose of describing constituent quarks.⁵ Rather, they were a more or less unavoidable step toward extracting the low-energy pseudoscalar degrees of freedom from the QCD Lagrangian.

If we compare the suggested gauge-fixed effective Lagrangian discussed above with the chiral quark model Lagrangian of eq.(110), we note two additional differences: The presence in our formulation of a Nakanishi-Lautrup field $b(x)$, and the ghosts $\bar{c}(x)$ and $c(x)$. These fields are of course all artifacts of the gauge-fixing formalism, and should be integrated out before a direct comparison can be made. The Faddeev-Popov ghosts, although presumably non-trivially coupled to the matrix $U(x)$ in the non-Abelian case, do not play a very fundamental rôle. Integrating them out produces an infinite series in derivatives of $U(x)$. These terms are of course crucial for, *e.g.*, all axial Ward identities to be satisfied in the rewritten theory, but we may think of them as only modifying the expansion implied in eq. (110).

A much more crucial rôle is played by the field $b(x)$. Integrating it out simply enforces the gauge-fixing constraint, and it is precisely by means of this constraint that one has the possibility of legitimately extracting the pion multiplet from the QCD Lagrangian. Of course, by means of exact rewritings of a generating functional for QCD, we do not learn about the preferred realization of chiral symmetry. Without an at least partial solution of essential QCD dynamics, we cannot see in what phase QCD may be realized in Nature. But the rewrite may still make the physical picture more transparent.

The point is the following. If we return to what would be our effective Lagrangian, then, in the absence of explicit chiral symmetry breaking in the form of current quark masses, we note that the pion multiplet appears only through derivative couplings. This is one of the standard features of chiral Lagrangians, implying the presence of massless pseudoscalars – the Goldstone bosons of the chiral symmetry breaking $SU(N_f)_L \times SU(N_f)_R \rightarrow SU(N_f)$. As we have stressed earlier, the collective fields introduced by the technique described in this paper are not tied to the possible existence of Goldstone modes. This is a question of dynamics. We may or may not have spontaneous symmetry breaking, and rewriting a generating functional of QCD adds nothing new to this. But *if* we end up extracting strictly massless pseudoscalars from the QCD Lagrangian, and if these fields are not trivially decoupled, then the left-over QCD-like theory of chirally rotated quark fields and gluons should be chirally non-invariant.⁶ This is basically the inverse of Goldstone’s Theorem.

Certainly such a lack of chiral invariance in the truncated part of our effective Lagrangian

⁵Collective fields corresponding explicitly to constituent quarks can presumably be introduced through suitable functional transformations as well. We have not pursued this idea, though. To do so, we would require a more precise definition of what is meant by constituent quarks in a Lagrangian framework. See below.

⁶We thank A. Manohar for emphasizing this point.

does not arise directly from an explicit constituent mass of the chirally rotated quarks, the $\bar{\chi}, \chi$ fields. If chiral symmetry is broken in the equivalent of the truncated theory, the possibility exists that it occurs as an explicit breaking due to the gauge-fixing constraint. This would be analogous to the Abelian case in two dimensions. For $SU(2)$ the gauge-fixing manifold would be a two-sphere. The gauge-fixing field $b(x)$ plays the rôle of a momentum conjugate to the angles describing the compact manifold. The global constraint on $b(x)$ is nothing else but the quantization condition of the corresponding angular momentum. Summation over the discrete values of the angular momentum, as in the Abelian case, projects on a certain point of the compact manifold and may thereby fix the values of the angles. Such a mechanism is possible in the $SU(N_f)$ -case too, and could imply chiral symmetry breaking in the truncated theory.

These considerations lead us to an interesting interpretation of the term “constituent quark”. In our framework one starts with current quarks. As one successively performs chiral rotations and the associated non-trivial gauge-fixings, one effectively removes the pseudoscalar degrees of freedom associated with the pairs of current quarks-antiquarks. If chiral symmetry is broken, then at each chiral rotation, the quarks become increasingly more “constituent”, and less “current”. This can be viewed as occurring through the fermion fields moving through the background of the pseudo-Goldstone fields $\theta(x)$. As there is never in this approach any double-counting of degrees of freedom, these chirally rotated quark fields are, in the phase of broken chiral symmetry, carrying less and less pseudoscalar degrees of freedom. In two space-time dimensions, where the same phenomenon should occur even in the abelian case, the chirally rotated quarks can almost entirely disappear in this process: This is the result of bosonization. It does not mean that constituent quarks cannot be given a meaningful definition after bosonization, but they now appear in the dual picture of a non-trivial soliton configuration [31]. In four dimensions, there is no possibility of constructing a topologically non-trivial configuration of the colour-singlet pseudoscalar degrees of freedom which carries the quantum numbers of a constituent quark. The chirally rotated quark fields $\chi, \bar{\chi}$ are then the fields we wish to identify with the constituent quarks. A related and perhaps complementary picture has been suggested by Kaplan [32], in which constituent quarks are viewed as Skyrmions in “colour space”. Also such an approach can be pursued in a collective-field framework similar to the one presented here for the flavoured case. Bosonization of QCD_2 indeed confirms this picture [31].

5 Conclusion

We have presented a new technique for the derivation of effective Lagrangians for long-distance dynamics, starting from an underlying Lagrangian valid to much smaller distances. The derivation relies heavily on a gauge-symmetric formulation that allows us to view different field representations of a given theory as nothing but different gauge slices.

As one application, this paper has focused on the classic problem of deriving an effective low-energy Lagrangian for the flavour-singlet sector of strong interactions, starting from QCD. The effective Lagrangian we have presented is equivalent (in the sense of generating functionals) to QCD with an explicit ultraviolet cut-off. It is connected directly to QCD through a series of well-defined field redefinitions in the cut-off theory, and in that sense is as fundamental as the underlying cut-off QCD Lagrangian itself. Technically, this is achieved by standard BRST quantization of the equivalent gauge-symmetric formulation.

The scheme is obviously valid in larger generality. We have already mentioned the case

of two-dimensional physics, where it provides an interpolation between purely fermionic and purely bosonic theories, thereby extending the meaning of bosonization in two dimensions [14]. We have also indicated how it can be extended to include the “pion” multiplets of $SU(N_f)$ as extracted from QCD. It can straightforwardly be applied to any Lagrangian from which one wishes to extract certain collective fields in an exact manner. Particular cases of interest may involve, as in QCD, a focus on Goldstone or pseudo-Goldstone fields described by coordinates on a quotient space G/H , corresponding to a general spontaneous symmetry-breaking pattern $G \rightarrow H$. Such a generalization may be of interest in the study of, *e.g.*, technicolour theories. But as we have repeatedly emphasized, collective fields may be of use much beyond the description of nearly-massless pseudo-Goldstone bosons.

If we return to the effective $U(1)$ Lagrangian derived above, where are the other pseudoscalars, the vector bosons, all the low-mass resonances, the baryons, etc. in this formalism? Since the effective Lagrangians we have provided examples of are exact rewrites of cut-off QCD, these other hadronic excitations below the cut-off are all included. We have not extracted them explicitly as collective fields, and they are therefore simply contained in the dynamics of the leftover (rotated) quark and gluon fields and their couplings to the explicit meson sector – just as also the lowest-mass pseudoscalars are implicitly contained in the starting QCD Lagrangian. If it is convenient to do so, one may introduce collective fields also for these higher excitations. This, incidentally, does not exclude the possibility that stable solitons with baryon quantum numbers can be constructed out of the collective pion fields. To the extent that the physical baryons indeed can be viewed as Skyrmions of meson multiplets, it is quite possible that extracting pion degrees of freedom from a generating functional of QCD also will entail a partial extraction of baryonic excitations. Following the rules laid out in this paper, there will by construction never be any double counting of the physical degrees of freedom.

References

- [1] B.W. Lee, “Chiral Dynamics”, Gordon and Breach Science Publ., New York (1972).
- [2] J. Gasser and H. Leutwyler, Phys. Lett. **B125** (1983) 321; Ann. Phys. (N.Y.) **158** (1984) 142; Nucl. Phys. **B250** (1985) 465, 517, 539.
- [3] G. Ecker, CERN preprint CERN-TH-6660/92 (1992).
J. Bijnens, Int. J. Mod. Phys. **A8** (1993) 3045.
- [4] K.G. Wilson and J. Kogut, Phys. Rep. **12** (1974) 75.
K.G. Wilson, Rev. Mod. Phys. **55** (1983) 583.
- [5] S. Weinberg, Physica **A96** (1979) 327.
- [6] H. Georgi, “Weak Interactions and Modern Particle Theory”, Benjamin/Cummings, Menlo Park (1984); Nucl. Phys. **B361** (1991) 339; Nucl. Phys. **B363** (1991) 301.
- [7] A. Manohar and H. Georgi, Nucl. Phys. **B234** (1984) 189.
- [8] G.P. Lepage, in “From Action to Answers” (T. DeGrand and D. Toussaint, Eds.) World Scientific Publ. Co., Singapore (1989).
J. Polchinski, preprint NSF-ITP-92-132.
- [9] R.I. Nepomechie, Ann. Phys. (N.Y.) **158** (1984) 67.
J. Balog, Phys. Lett. **B149** (1984) 197.
A.A. Andrianov and L. Bonora, Nucl. Phys. **B233** (1984) 232.
A.A. Andrianov, Phys. Lett. **B157** (1985) 425.
R.D. Ball and H. Osborn, Phys. Lett. **B165** (1985) 410.
I.J.R. Aitchison and C.M. Fraser, Phys. Rev. **D31** (1985) 2605.
D.W. McKay and M.J. Munczek, Phys. Rev. **D32** (1985) 266.
P. Simic, Phys. Rev. Lett. **55** (1985) 40; Phys. Rev. **D34** (1986) 1903.
J. Zuk, Z. Phys. **C29** (1985) 303.
- [10] D. Espriu, R. de Rafael and J. Taron, Nucl. Phys. **B345** (1990) 22.
- [11] Proceedings of the Workshop on Skyrmions and Anomalies (M. Jezabek and M. Praszalowicz, Eds.), World Scientific Publ. Co., Singapore (1987).
- [12] A. Zaks, Nucl. Phys. **B260** (1985) 241.
- [13] N.I. Karchev and A.A. Slavnov, Theor. Math. Phys. **65** (1986) 1099.
- [14] P.H. Damgaard, H.B. Nielsen and R. Sollacher, Nucl. Phys. **B385** (1992) 227; Phys. Lett. **B296** (1992) 132.
- [15] T. Ikehashi, preprint UT-Komaba 93-11, hep-th/9307136.
- [16] E. Witten, Nucl. Phys. **B156** (1979) 269.
- [17] G. Veneziano, Nucl. Phys. **B159** (1979) 213.

- [18] J. Schwinger, Phys. Rev. **128** (1962) 2425;
J. Lowenstein and J. Swieca, Ann. Phys. (N.Y.) **68** (1971) 175.
- [19] G. Morchio, D. Pierotti and F. Strocchi, Ann. Phys. (N.Y.) **188** (1988) 217.
- [20] A.V. Manohar, Phys. Rev. Lett. **66** (1991) 1663.
- [21] R.D. Ball, Phys. Rep. **182** (1989) 1.
- [22] A. Hosoya and K. Kikkawa, Nucl. Phys. **B101** (1975) 271.
- [23] J. Alfaro and P.H. Damgaard, Ann. Phys. (NY) **202** (1990) 398.
For an earlier discussion, see also D. Förster, H.B. Nielsen and M. Ninomiya, Phys. Lett. **B94** (1980) 135.
- [24] C. Jayewardena, Helv. Phys. Acta **61** (1988) 636.
I. Sachs and A. Wipf, preprint ETH-TH/91-15.
- [25] N.A. McDougall, Nucl. Phys. **B211** (1983) 139.
R.D. Carlitz, Nucl. Phys. **B236** (1984) 423.
N.J. Dowrick and N.A. McDougall, Phys. Lett. **B285** (1992) 269.
- [26] G.M. Shore, Nucl. Phys. **B362** (1991) 85.
- [27] G.M. Shore and G. Veneziano, Nucl. Phys. **B381** (1992) 3.
- [28] J. Polchinski, Nucl. Phys. **B231** (1984) 269.
B.J. Warr, Ann. Phys.(N.Y.) **183** (1988) 1, 59.
- [29] K. Bardakci and L. Crescimanno, Nucl. Phys. **B313** (1989) 269.
M.V. Manias, C.M. Naon and M.L. Trobo, Phys. Rev. **D41** (1990) 3174.
- [30] J. Kurchan, D.R. Bes and S. Cruz Barrios, Phys. Rev. **D38** (1988) 3309.
- [31] J. Ellis, Y. Frishman and M. Karliner, Phys. Lett. **B272** (1991) 333.
J. Ellis, Y. Frishman, A. Hanany and M. Karliner, Nucl. Phys. **B382** (1992) 189.
- [32] D. Kaplan, Phys. Lett. **B235** (1990) 163; Nucl. Phys. **B351** (1991) 137.